

## *Testing $B$ -mesogenesis at LHCb*

Alexandre Brea Rodríguez, Veronika Chobanova<sup>1</sup>, Xabier Cid Vidal,  
Saúl López Soliño, Titus Mombacher, Claire Prouvé, Emilio Xosé Rodríguez Fernández  
(USC)

Diego Martínez Santos (GAIN/USC)

Carlos Vázquez Sierra (CERN)


[\[arXiv:2106.12870\]](https://arxiv.org/abs/2106.12870)

<sup>1</sup>On maternity leave




# Introduction

B- mesogenesis: Alonso-Álvarez, Escudero, and Elor [\[arXiv:2101.02706\]](https://arxiv.org/abs/2101.02706)





## Introduction



Gonzalo Alonso-Álvarez

Similar spirit to Hylogenesis,  
Davoudiasl et al. [\[arXiv:1008.2399\]](https://arxiv.org/abs/1008.2399)

$$Y_{\mathcal{B}} \simeq 8.7 \times 10^{-11} \frac{\text{Br}(B \rightarrow \psi + \mathcal{B} + \mathcal{M})}{10^{-2}} \sum_{q=s,d} \alpha_q \frac{A_{\text{SL}}^q}{10^{-4}}$$




# Ingredients for baryon asymmetry

$$Y_B \propto \sum_{s,d} A_{SL}^{s,d} \times \mathcal{B}(B_{s,d}^0 \rightarrow \psi_{DS} X)$$

At least one of the two  $A_{sl}$  needs to be **positive** to produce baryon abundance

Indirect determinations significantly favor a negative value for  $B_d$ , and in SM at least a **positive value for  $B_s$** .

Directly measured at LHCb and other experiments



$$B_d : A_{SL} = -0.0021 \pm 0.0017$$

$$B_s : A_{SL} = -0.0006 \pm 0.0028$$



## Ingredients for baryon asymmetry

$$Y_B \propto \sum_{s,d} A_{SL}^{s,d} \times \mathcal{B}(B_{s,d}^0 \rightarrow \psi_{DS} X)$$

At least one of the two  $A_{sl}$  needs to be **positive** to produce baryon abundance

Indirect determinations significantly favor a negative value for  $B_d$ , and in SM at least a **positive value for  $B_s$** .

Indirect determinations are in principle more precise

$$A_{SL}^d |_{SM} = (-4.7 \pm 0.4) \times 10^{-4}$$

$$A_{SL}^s |_{SM} = (2.1 \pm 0.2) \times 10^{-5}$$

Lenz & Tetlalmatzi-Xolocotzi  
1912.07621

UTfit  
NP-fit  
2018

$A_{SL_d}$	$-0.0033 \pm 0.0014$
$A_{SL_s}$	$-0.00013 \pm 0.00051$

→  $B_s$  is the likeliest candidate for B-mesogenesis



## Ingredients for baryon asymmetry

$$Y_B \propto \sum_{s,d} A_{SL}^{s,d} \times \mathcal{B}(B_{s,d}^0 \rightarrow \psi_{DS} X) =$$

$$- \sum_{s,d} \left| \frac{\Delta \Gamma_{s,d}}{\Delta m_{s,d} \cos(\phi_{s,d} - \arg(\Gamma_{12}^{s,d}))} \right| \sin(\phi_{s,d} - \arg(\Gamma_{12}^{s,d}))$$

$$\times \mathcal{B}(B_{s,d}^0 \rightarrow \psi_{DS} X) ,$$

Calculated in global fits (e.g. UTfit)

Very precise ( $\sim 2$  mrad) in SM and models w/o significant BSM contributions to  $\Delta F = 1$  penguins

**Some theory work needed** to compute it in the context of B-mesogenesis (e.g., whether it adds BSM contributions to  $\Delta F = 1$  penguins)



## Ingredients for baryon asymmetry


$$Y_B \propto \sum_{s,d} A_{SL}^{s,d} \times \mathcal{B}(B_{s,d}^0 \rightarrow \psi_{DS} X) =$$

$$- \sum_{s,d} \left| \frac{\Delta \Gamma_{s,d}}{\Delta m_{s,d} \cos(\phi_{s,d} - \arg(\Gamma_{12}^{s,d}))} \right| \sin(\phi_{s,d} - \arg(\Gamma_{12}^{s,d}))$$

$$\times \mathcal{B}(B_{s,d}^0 \rightarrow \psi_{DS} X) ,$$

LHCb

—  $B_s^0 \rightarrow D_s^- \pi^+$  —  $\bar{B}_s^0 \rightarrow D_s^- \pi^+$  — Untagged



[\[arXiv:2104.04421\]](https://arxiv.org/abs/2104.04421)

$$\Delta m_s = 17.7656 \pm 0.0057 \text{ ps}^{-1}$$




## Ingredients for baryon asymmetry

LHCb (& ATLAS/CMS)

$$Y_B \propto \sum_{s,d} A_{SL}^{s,d} \times \mathcal{B}(B_{s,d}^0 \rightarrow \psi_{DS} X) =$$

$$- \sum_{s,d} \left| \frac{\Delta \Gamma_{s,d}}{\Delta m_{s,d} \cos(\phi_{s,d} - \arg(\Gamma_{12}^{s,d}))} \right| \sin(\phi_{s,d} - \arg(\Gamma_{12}^{s,d}))$$

$$\times \mathcal{B}(B_{s,d}^0 \rightarrow \psi_{DS} X) ,$$






## *Ingredients for baryon asymmetry*

**LHCb ?** → this talk

$$\begin{aligned}
 Y_B &\propto \sum_{s,d} A_{SL}^{s,d} \times \mathcal{B}(B_{s,d}^0 \rightarrow \psi_{DS} X) = \\
 &- \sum_{s,d} \left| \frac{\Delta \Gamma_{s,d}}{\Delta m_{s,d} \cos(\phi_{s,d} - \arg(\Gamma_{12}^{s,d}))} \right| \sin(\phi_{s,d} - \arg(\Gamma_{12}^{s,d})) \\
 &\times \mathcal{B}(B_{s,d}^0 \rightarrow \psi_{DS} X)
 \end{aligned}$$





## Setup






## Setup: Fast simulation

- + Fast simulation of the tracking system, including multiple scattering (V. Chobanova et al. [2012.02692](#))
- + Particle identification from existing plots in LHCb papers/TDR's







# Signatures



The b-quark decays, the other(s) are spectator  
 → can be searched using **any of the b-hadron species,**  
**no need to restrict to Bs**

Inclusive BR's:  $\sim 0.1 - 0.001$


**Exclusive BR's:  $\sim 0.1 - 10^{-6}$**






## Signatures

No tracks nearby



Daughter tracks close to each other (DOCA)



Usual requirements for b-daughter tracks:  $P_T$ , detachment from PV.



# Efficiencies

Decay mode	$\epsilon^{REC}\%$	$\epsilon^{REC\&PT}$	$\epsilon^{SEL/REC}$	$\epsilon^{PID/SEL}$
$\Lambda_b^0 \rightarrow \psi_{DS}(940)K^+\pi^-$	7.6	5.1	13.2	74.2
$\Lambda_b^0 \rightarrow \psi_{DS}(940)\pi^+\pi^-$	7.3	4.8	14.3	-
$\Lambda_b^0 \rightarrow \psi_{DS}(1500)K^+\pi^-$	7.7	4.8	10.8	76.0
$\Lambda_b^0 \rightarrow \psi_{DS}(1500)\pi^+\pi^-$	7.3	4.5	12.5	-
$\Lambda_b^0 \rightarrow \psi_{DS}(2000)K^+\pi^-$	7.7	4.5	6.67	79.2
$\Lambda_b^0 \rightarrow \psi_{DS}(2000)\pi^+\pi^-$	7.4	4.1	10.9	-
$\Lambda_b^0 \rightarrow \psi_{DS}(2400)K^+\pi^-$	7.8	4.1	9.11	80.8
$\Lambda_b^0 \rightarrow \psi_{DS}(2400)\pi^+\pi^-$	7.4	3.7	8.89	-
$\Lambda_b^0 \rightarrow \psi_{DS}(4340)K^+\pi^-$	8.1	1.2	3.12	88.0
$\Lambda_b^0 \rightarrow \psi_{DS}(4340)\pi^+\pi^-$	7.4	0.97	2.88	-
$\Lambda_b^0 \rightarrow \psi_{DS}(4470)K^+\pi^-$	8.2	0.91	2.18	87.7
$\Lambda_b^0 \rightarrow \psi_{DS}(4470)\pi^+\pi^-$	7.4	0.79	2.23	-
$B^+ \rightarrow \psi_{DS}(940)\Lambda_c(2595)^+$	5.7	1.3	20.9	56.7
$B^+ \rightarrow \psi_{DS}(1500)\Lambda_c(2595)^+$	5.6	1.1	19.3	56.6
$B^+ \rightarrow \psi_{DS}(2000)\Lambda_c(2595)^+$	5.3	0.88	18.5	57.2
$B^+ \rightarrow \psi_{DS}(2400)\Lambda_c(2595)^+$	4.9	0.66	16.4	57.1
$B^0 \rightarrow \psi_{DS}(940)\Lambda(1520)$	13.3	12.7	38.1	56.9
$B^0 \rightarrow \psi_{DS}(1500)\Lambda(1520)$	13.7	12.9	36.6	56.4
$B^0 \rightarrow \psi_{DS}(2000)\Lambda(1520)$	13.5	12.6	35.5	56.8
$B^0 \rightarrow \psi_{DS}(2400)\Lambda(1520)$	13.3	6.8	34.5	56.6
$B^0 \rightarrow \psi_{DS}(3500)\Lambda(1520)$	12.1	3.5	23.6	56.5

Used 4 signal decays

$$\Lambda_b \rightarrow \psi_{DS} K \pi$$

$$\Lambda_b \rightarrow \psi_{DS} \pi \pi$$

$$B_d \rightarrow \psi_{DS} \Lambda(1520)$$

$$B^+ \rightarrow \psi_{DS} \Lambda_c(2595)^+$$



# Efficiencies

Decay mode	$\epsilon^{REC}\%$	$\epsilon^{REC\&PT}$	$\epsilon^{SEL/REC}$	$\epsilon^{PID/SEL}$
$\Lambda_b^0 \rightarrow \psi_{DS}(940)K^+\pi^-$	7.6	5.1	13.2	74.2
$\Lambda_b^0 \rightarrow \psi_{DS}(940)\pi^+\pi^-$	7.3	4.8	14.3	-
$\Lambda_b^0 \rightarrow \psi_{DS}(1500)K^+\pi^-$	7.7	4.8	10.8	76.0
$\Lambda_b^0 \rightarrow \psi_{DS}(1500)\pi^+\pi^-$	7.3	4.5	12.5	-
$\Lambda_b^0 \rightarrow \psi_{DS}(2000)K^+\pi^-$	7.7	4.5	6.67	79.2
$\Lambda_b^0 \rightarrow \psi_{DS}(2000)\pi^+\pi^-$	7.4	4.1	10.9	-
$\Lambda_b^0 \rightarrow \psi_{DS}(2400)K^+\pi^-$	7.8	4.1	9.11	80.8
$\Lambda_b^0 \rightarrow \psi_{DS}(2400)\pi^+\pi^-$	7.4	3.7	8.89	-
$\Lambda_b^0 \rightarrow \psi_{DS}(4340)K^+\pi^-$	8.1	1.2	3.12	88.0
$\Lambda_b^0 \rightarrow \psi_{DS}(4340)\pi^+\pi^-$	7.4	0.97	2.88	-
$\Lambda_b^0 \rightarrow \psi_{DS}(4470)K^+\pi^-$	8.2	0.91	2.18	87.7
$\Lambda_b^0 \rightarrow \psi_{DS}(4470)\pi^+\pi^-$	7.4	0.79	2.23	-
$B^+ \rightarrow \psi_{DS}(940)\Lambda_c(2595)^+$	5.7	1.3	20.9	56.7
$B^+ \rightarrow \psi_{DS}(1500)\Lambda_c(2595)^+$	5.6	1.1	19.3	56.6
$B^+ \rightarrow \psi_{DS}(2000)\Lambda_c(2595)^+$	5.3	0.88	18.5	57.2
$B^+ \rightarrow \psi_{DS}(2400)\Lambda_c(2595)^+$	4.9	0.66	16.4	57.1
$B^0 \rightarrow \psi_{DS}(940)\Lambda(1520)$	13.3	12.7	38.1	56.9
$B^0 \rightarrow \psi_{DS}(1500)\Lambda(1520)$	13.7	12.9	36.6	56.4
$B^0 \rightarrow \psi_{DS}(2000)\Lambda(1520)$	13.5	12.6	35.5	56.8
$B^0 \rightarrow \psi_{DS}(2400)\Lambda(1520)$	13.3	6.8	34.5	56.6
$B^0 \rightarrow \psi_{DS}(3500)\Lambda(1520)$	12.1	3.5	23.6	56.5

Reconstruction efficiency  
(including detector  
acceptance), in %



## Efficiencies

Decay mode	$\epsilon^{REC}\%$	$\epsilon^{REC\&PT}$	$\epsilon^{SEL/REC}$	$\epsilon^{PID/SEL}$
$\Lambda_b^0 \rightarrow \psi_{DS}(940)K^+\pi^-$	7.6	5.1	13.2	74.2
$\Lambda_b^0 \rightarrow \psi_{DS}(940)\pi^+\pi^-$	7.3	4.8	14.3	-
$\Lambda_b^0 \rightarrow \psi_{DS}(1500)K^+\pi^-$	7.7	4.8	10.8	76.0
$\Lambda_b^0 \rightarrow \psi_{DS}(1500)\pi^+\pi^-$	7.3	4.5	12.5	-
$\Lambda_b^0 \rightarrow \psi_{DS}(2000)K^+\pi^-$	7.7	4.5	6.67	79.2
$\Lambda_b^0 \rightarrow \psi_{DS}(2000)\pi^+\pi^-$	7.4	4.1	10.9	-
$\Lambda_b^0 \rightarrow \psi_{DS}(2400)K^+\pi^-$	7.8	4.1	9.11	80.8
$\Lambda_b^0 \rightarrow \psi_{DS}(2400)\pi^+\pi^-$	7.4	3.7	8.89	-
$\Lambda_b^0 \rightarrow \psi_{DS}(4340)K^+\pi^-$	8.1	1.2	3.12	88.0
$\Lambda_b^0 \rightarrow \psi_{DS}(4340)\pi^+\pi^-$	7.4	0.97	2.88	-
$\Lambda_b^0 \rightarrow \psi_{DS}(4470)K^+\pi^-$	8.2	0.91	2.18	87.7
$\Lambda_b^0 \rightarrow \psi_{DS}(4470)\pi^+\pi^-$	7.4	0.79	2.23	-
$B^+ \rightarrow \psi_{DS}(940)\Lambda_c(2595)^+$	5.7	1.3	20.9	56.7
$B^+ \rightarrow \psi_{DS}(1500)\Lambda_c(2595)^+$	5.6	1.1	19.3	56.6
$B^+ \rightarrow \psi_{DS}(2000)\Lambda_c(2595)^+$	5.3	0.88	18.5	57.2
$B^+ \rightarrow \psi_{DS}(2400)\Lambda_c(2595)^+$	4.9	0.66	16.4	57.1
$B^0 \rightarrow \psi_{DS}(940)\Lambda(1520)$	13.3	12.7	38.1	56.9
$B^0 \rightarrow \psi_{DS}(1500)\Lambda(1520)$	13.7	12.9	36.6	56.4
$B^0 \rightarrow \psi_{DS}(2000)\Lambda(1520)$	13.5	12.6	35.5	56.8
$B^0 \rightarrow \psi_{DS}(2400)\Lambda(1520)$	13.3	6.8	34.5	56.6
$B^0 \rightarrow \psi_{DS}(3500)\Lambda(1520)$	12.1	3.5	23.6	56.5

Reconstruction efficiency  
(including detector acceptance),  
in %, & 2 tracks with PT > 800 MeV



# Efficiencies

Decay mode	$\epsilon^{REC}\%$	$\epsilon^{REC\&PT}$	$\epsilon^{SEL/REC}$	$\epsilon^{PID/SEL}$
$\Lambda_b^0 \rightarrow \psi_{DS}(940)K^+\pi^-$	7.6	5.1	13.2	74.2
$\Lambda_b^0 \rightarrow \psi_{DS}(940)\pi^+\pi^-$	7.3	4.8	14.3	-
$\Lambda_b^0 \rightarrow \psi_{DS}(1500)K^+\pi^-$	7.7	4.8	10.8	76.0
$\Lambda_b^0 \rightarrow \psi_{DS}(1500)\pi^+\pi^-$	7.3	4.5	12.5	-
$\Lambda_b^0 \rightarrow \psi_{DS}(2000)K^+\pi^-$	7.7	4.5	6.67	79.2
$\Lambda_b^0 \rightarrow \psi_{DS}(2000)\pi^+\pi^-$	7.4	4.1	10.9	-
$\Lambda_b^0 \rightarrow \psi_{DS}(2400)K^+\pi^-$	7.8	4.1	9.11	80.8
$\Lambda_b^0 \rightarrow \psi_{DS}(2400)\pi^+\pi^-$	7.4	3.7	8.89	-
$\Lambda_b^0 \rightarrow \psi_{DS}(4340)K^+\pi^-$	8.1	1.2	3.12	88.0
$\Lambda_b^0 \rightarrow \psi_{DS}(4340)\pi^+\pi^-$	7.4	0.97	2.88	-
$\Lambda_b^0 \rightarrow \psi_{DS}(4470)K^+\pi^-$	8.2	0.91	2.18	87.7
$\Lambda_b^0 \rightarrow \psi_{DS}(4470)\pi^+\pi^-$	7.4	0.79	2.23	-
$B^+ \rightarrow \psi_{DS}(940)\Lambda_c(2595)^+$	5.7	1.3	20.9	56.7
$B^+ \rightarrow \psi_{DS}(1500)\Lambda_c(2595)^+$	5.6	1.1	19.3	56.6
$B^+ \rightarrow \psi_{DS}(2000)\Lambda_c(2595)^+$	5.3	0.88	18.5	57.2
$B^+ \rightarrow \psi_{DS}(2400)\Lambda_c(2595)^+$	4.9	0.66	16.4	57.1
$B^0 \rightarrow \psi_{DS}(940)\Lambda(1520)$	13.3	12.7	38.1	56.9
$B^0 \rightarrow \psi_{DS}(1500)\Lambda(1520)$	13.7	12.9	36.6	56.4
$B^0 \rightarrow \psi_{DS}(2000)\Lambda(1520)$	13.5	12.6	35.5	56.8
$B^0 \rightarrow \psi_{DS}(2400)\Lambda(1520)$	13.3	6.8	34.5	56.6
$B^0 \rightarrow \psi_{DS}(3500)\Lambda(1520)$	12.1	3.5	23.6	56.5

Reconstruction efficiency  
(including detector acceptance),  
in %, & 2 tracks with PT > 800 MeV



# Efficiencies

Decay mode	$\epsilon^{REC}\%$	$\epsilon^{REC\&PT}$	$\epsilon^{SEL/REC}$	$\epsilon^{PID/SEL}$
$\Lambda_b^0 \rightarrow \psi_{DS}(940)K^+\pi^-$	7.6	5.1	13.2	74.2
$\Lambda_b^0 \rightarrow \psi_{DS}(940)\pi^+\pi^-$	7.3	4.8	14.3	-
$\Lambda_b^0 \rightarrow \psi_{DS}(1500)K^+\pi^-$	7.7	4.8	10.8	76.0
$\Lambda_b^0 \rightarrow \psi_{DS}(1500)\pi^+\pi^-$	7.3	4.5	12.5	-
$\Lambda_b^0 \rightarrow \psi_{DS}(2000)K^+\pi^-$	7.7	4.5	6.67	79.2
$\Lambda_b^0 \rightarrow \psi_{DS}(2000)\pi^+\pi^-$	7.4	4.1	10.9	-
$\Lambda_b^0 \rightarrow \psi_{DS}(2400)K^+\pi^-$	7.8	4.1	9.11	80.8
$\Lambda_b^0 \rightarrow \psi_{DS}(2400)\pi^+\pi^-$	7.4	3.7	8.89	-
$\Lambda_b^0 \rightarrow \psi_{DS}(4340)K^+\pi^-$	8.1	1.2	3.12	88.0
$\Lambda_b^0 \rightarrow \psi_{DS}(4340)\pi^+\pi^-$	7.4	0.97	2.88	-
$\Lambda_b^0 \rightarrow \psi_{DS}(4470)K^+\pi^-$	8.2	0.91	2.18	87.7
$\Lambda_b^0 \rightarrow \psi_{DS}(4470)\pi^+\pi^-$	7.4	0.79	2.23	-
$B^+ \rightarrow \psi_{DS}(940)\Lambda_c(2595)^+$	5.7	1.3	20.9	56.7
$B^+ \rightarrow \psi_{DS}(1500)\Lambda_c(2595)^+$	5.6	1.1	19.3	56.6
$B^+ \rightarrow \psi_{DS}(2000)\Lambda_c(2595)^+$	5.3	0.88	18.5	57.2
$B^+ \rightarrow \psi_{DS}(2400)\Lambda_c(2595)^+$	4.9	0.66	16.4	57.1
$B^0 \rightarrow \psi_{DS}(940)\Lambda(1520)$	13.3	12.7	38.1	56.9
$B^0 \rightarrow \psi_{DS}(1500)\Lambda(1520)$	13.7	12.9	36.6	56.4
$B^0 \rightarrow \psi_{DS}(2000)\Lambda(1520)$	13.5	12.6	35.5	56.8
$B^0 \rightarrow \psi_{DS}(2400)\Lambda(1520)$	13.3	6.8	34.5	56.6
$B^0 \rightarrow \psi_{DS}(3500)\Lambda(1520)$	12.1	3.5	23.6	56.5

Selection efficiency for those reconstructed events.

Selection includes PT cuts, vertex quality, isolation...


Not yet PT miss or invariant mass



# Efficiencies



Decay mode	$\epsilon^{REC}\%$	$\epsilon^{REC\&PT}$	$\epsilon^{SEL/REC}$	$\epsilon^{PID/SEL}$
$\Lambda_b^0 \rightarrow \psi_{DS}(940)K^+\pi^-$	7.6	5.1	13.2	74.2
$\Lambda_b^0 \rightarrow \psi_{DS}(940)\pi^+\pi^-$	7.3	4.8	14.3	-
$\Lambda_b^0 \rightarrow \psi_{DS}(1500)K^+\pi^-$	7.7	4.8	10.8	76.0
$\Lambda_b^0 \rightarrow \psi_{DS}(1500)\pi^+\pi^-$	7.3	4.5	12.5	-
$\Lambda_b^0 \rightarrow \psi_{DS}(2000)K^+\pi^-$	7.7	4.5	6.67	79.2
$\Lambda_b^0 \rightarrow \psi_{DS}(2000)\pi^+\pi^-$	7.4	4.1	10.9	-
$\Lambda_b^0 \rightarrow \psi_{DS}(2400)K^+\pi^-$	7.8	4.1	9.11	80.8
$\Lambda_b^0 \rightarrow \psi_{DS}(2400)\pi^+\pi^-$	7.4	3.7	8.89	-
$\Lambda_b^0 \rightarrow \psi_{DS}(4340)K^+\pi^-$	8.1	1.2	3.12	88.0
$\Lambda_b^0 \rightarrow \psi_{DS}(4340)\pi^+\pi^-$	7.4	0.97	2.88	-
$\Lambda_b^0 \rightarrow \psi_{DS}(4470)K^+\pi^-$	8.2	0.91	2.18	87.7
$\Lambda_b^0 \rightarrow \psi_{DS}(4470)\pi^+\pi^-$	7.4	0.79	2.23	-
$B^+ \rightarrow \psi_{DS}(940)\Lambda_c(2595)^+$	5.7	1.3	20.9	56.7
$B^+ \rightarrow \psi_{DS}(1500)\Lambda_c(2595)^+$	5.6	1.1	19.3	56.6
$B^+ \rightarrow \psi_{DS}(2000)\Lambda_c(2595)^+$	5.3	0.88	18.5	57.2
$B^+ \rightarrow \psi_{DS}(2400)\Lambda_c(2595)^+$	4.9	0.66	16.4	57.1
$B^0 \rightarrow \psi_{DS}(940)\Lambda(1520)$	13.3	12.7	38.1	56.9
$B^0 \rightarrow \psi_{DS}(1500)\Lambda(1520)$	13.7	12.9	36.6	56.4
$B^0 \rightarrow \psi_{DS}(2000)\Lambda(1520)$	13.5	12.6	35.5	56.8
$B^0 \rightarrow \psi_{DS}(2400)\Lambda(1520)$	13.3	6.8	34.5	56.6
$B^0 \rightarrow \psi_{DS}(3500)\Lambda(1520)$	12.1	3.5	23.6	56.5

Particle Identification efficiency  
for selected events  
(Separation of Kaon/proton vs  
pion)







$$B_d \rightarrow \psi_{DS} A(1520), B^+ \rightarrow \psi_{DS} A_c(2595)^+$$



→ Search for signal in bins of  $p_T^{\text{miss}}$  assuming BKG level is known



$$\Lambda_b \rightarrow \psi_{DS} K\pi, \Lambda_b \rightarrow \psi_{DS} \pi\pi$$




- MVA combining  $M(h,h)$  &  $PT_{\text{miss}}$
- Optimize cut and search for signal assuming BKG level is known




# Sensitivity


LHCb Upgrade  
(Phase-I)



$50 \text{ fb}^{-1}$




LHCb Upgrade  
(Phase-II)





## Sensitivity




→ Signals are cleaner at low masses of the  $\psi_{DS}$  for the same BR → stronger BR limits at low masses

→ On the other hand, if  $\psi_{DS}$  is heavy, the exclusive BR's are higher → no need to go to so low BR to probe the theory

→ In terms of sample size, LHCb Upgrade can probe all allowed mass range for  $\psi_{DS}$



## Comment on systematics



A crucial difference w.r.t other searches is the extremely high background level

→ Even with small (per mil) systematics the measurement would be systematically limited after 10 fb<sup>-1</sup>



→ In terms of sample size, LHCb Upgrade can probe all allowed mass range for  $\psi_{DS}$

→ LHCb Upgrade needs very precise bkg systematic (and/or stronger bkg suppression) to probe all allowed mass range for  $\psi_{DS}$



## Use of mother resonance?

Further bkg suppression (as well as a clean signal peak confirmation) can be achieved by identifying the mother of the decaying b-hadron (e.g.,  $\Sigma_b^+ \rightarrow \Lambda_b \pi^+$ ) (see Zhang&Stone, [1402.4205](#))





## Conclusions

- B-mesogenesis is an elegant solution that solves the baryon asymmetry problem by eliminating the baryon asymmetry
- It can be accurately tested by precise measurements of  $B_{s(d)}$  oscillation parameters and searches for b decays to invisible + X
- LHCb can search for those BSM decays with excellent **statistical** accuracy
- Though **systematics** from background modelling **could be a limiting factor**
- Further bkg suppression (as well as a clean signal peak confirmation) can be achieved by identifying the mother of the decaying b-handron (eg ,  $\Sigma_b^+ \rightarrow \Lambda_b \pi^+$ )
- LHCb has the **potential** to test the entire parameter space of the model



*Backup*



