


A short review of spin physics



Barbara Badelek
University of Warsaw

Low x 2010

Kavala, June 23 – 27, 2010

A Beautiful Spin (after X. Ji)

- Born with troubles (Stern & Gerlach (1922) vs Goudsmit & Uhlenbeck (1925))
- Is due to space–time symmetry
- Fundamental concept
- **Laboratory to explore physics beyond the SM, e.g.:**
 - Muon “ $g - 2$ ” experiment @ BNL
 - Proton weak charge (Qweak exp @ JLAB)
 - Neutron EDM measurement ...


Tool to measure observables hard to obtain otherwise, e.g:

- Strangeness content of the nucleon from polarised parity–violating e–p scattering
- Electromagnetic form factors of the nucleon from the recoil polarisation
- Neutron density in large nuclei from parity–violating electron scattering
- and...

A Beautiful Spin (after X. Ji)...cont'd

- Probe to unravel the nonperturbative QCD dynamics, e.g.:
 - Nucleon spin-dependent structure functions, g_1 and g_2
 - Quark helicity ($\Delta q(x)$) and transversity ($\Delta_T q(x)$) distributions
 - Gluon polarisation, $\Delta g(x)$
 - Generalised Parton Distributions, GPD
 - Semi-Inclusive Deep Inelastic Scattering, SIDIS
 - (Generalised) Drell-Hearn-Gerasimov-... sum rule
 - Single spin asymmetries


Nucleon spin structure in the electroproduction




- $\frac{d^2\sigma}{d\Omega dE'} = \frac{\alpha^2}{2Mq^4} \frac{E'}{E} L_{\mu\nu} W^{\mu\nu}$
- Symmetric part of $W^{\mu\nu}$ – unpol. DIS, antisymmetric – polarised DIS
- Nominally $F_{1,2}$, $q(x) \longrightarrow g_{1,2}$, $\Delta q(x)$ but...
- ...anomalous gluon contribution to $g_1(x)$
- ... $g_2(x)$ has no interpretation in terms of partons.

Partonic structure of the nucleon; distribution functions


Three **twist-two** quark distributions in QCD (after integrating over the quark intrinsic k_t)



Quark momentum DF;
well known (unpolarised DIS $\rightarrow F_{1,2}(x)$).



Difference in DF of quarks with spin parallel or antiparallel to the nucleon's spin;
known (polarised DIS $\rightarrow g_1(x)$).



Difference in DF of quarks with spin parallel or antiparallel to the nucleon's spin in the transversely polarised nucleon;
unknown (polarised DIS $\rightarrow h_1(x)$).

Nonrelativistically: $\Delta_T q(x) \equiv \Delta q(x)$. **OBS.!** $\Delta_T q(x)$ are **C-odd and chiral-odd**; may only be measured with another chiral-odd partner, e.g. fragmentation function.

If the k_t taken into account \implies 8 TMD distr.; one, f_{1T}^\perp accessible through "Sivers asymmetry".

Nucleon spin structure: observables in $\vec{\mu}\vec{N}$ scattering

- Inclusive asymmetry, A_{meas} :

$$A_{meas} = \frac{1}{fP_T P_B} \left(\frac{N^{\leftrightarrow} - N^{\equiv}}{N^{\leftrightarrow} + N^{\equiv}} \right) \approx DA_1 = D \frac{g_1(x, Q^2)}{F_1(x, Q^2)} = D \frac{\sum_q e_q^2 \Delta q(x, Q^2)}{\sum_q e_q^2 q(x, Q^2)}$$


$$\Delta q = q^+ - q^-, \quad q = q^+ + q^-, \quad g_1^d = g_1^N \left(1 - \frac{3}{2}\omega_D\right) = \frac{g_1^p + g_1^n}{2} \left(1 - \frac{3}{2}\omega_D\right);$$

$$\omega_D = 0.05 \pm 0.01$$

- Semi-inclusive asymmetry, A_1^h :



$$A_1^h(x, z, Q^2) \approx D \frac{\sum_q e_q^2 \Delta q(x, Q^2) D_q^h(z, Q^2)}{\sum_q e_q^2 q(x, Q^2) D_q^h(z, Q^2)} \quad z = \frac{E_h}{\nu} \quad D_q^h \neq D_{\bar{q}}^h$$

Experiments




Experiment	Polarised beam	Polarised target	Energy (GeV)
SLAC	e	p, n, d	$\lesssim 50$
EMC	μ	p	100–200
SMC	μ	p, d	100, 190
HERMES	e	p, n, d	27.5
COMPASS	μ	p, d	160
JLAB	e	p, n, d	$\lesssim 6$

COMPASS polarised targets



- * Two (three in 2006, 2007) target cells, oppositely polarised
- * Polarisation reversed every 8 h (less frequent in 2006, 2007) by field rotation
- * Material: solid ${}^6\text{LiD}$ (NH_3 in 2007)
- * Polarisation: $\sim 50\%$ ($\sim 90\%$ in 2007), by the Dynamical Nuclear Polarisation
- * Dilution: $f \sim 0.4$ (~ 0.15 in 2007)
- * Polar acceptance: ~ 70 mrad (~ 180 mrad in 2006, 2007)

Polarised pp scattering at RHIC



STAR sees 4 helicity configurations
STAR runs 4 parallel measurements


RHIC measured polarization
Run 9 @ 2x250 GeV

Pol yellow 0.40

Pol blue 0.38

syst. pol (blue+yellow)=9.2%

$g_1(x)$ for proton and deuteron, $Q^2 > 1$ (GeV/c)²



Very precise data
especially at $x \lesssim 0.01$

From Γ_1^d at $Q^2 \rightarrow \infty$:

HERMES:


$$a_0 = 0.330 \pm 0.025 \pm 0.011 \pm 0.028$$

COMPASS:



$$a_0 = 0.33 \pm 0.03 \pm 0.05$$

$$\Delta s + \Delta \bar{s} = -0.08 \pm 0.01 \pm 0.02$$

Good agreement
between the experiments





$g_1(x)$ for proton and deuteron, $Q^2 > 1$ (GeV/c)²...cont'd



More data from JLAB/EG1a @ low Q^2 .




$g_1^d(x)$ in the nonperturbative ($Q^2 < 1$ (GeV/c) 2 region)

V.Yu. Alexakhin *et al.* (COMPASS) Phys. Lett. B **647** (2007) 330



- Order of magnitude improvement over the statistical precision of the SMC.
- Interplay between perturbative and nonperturbative mechanisms.
- Spin effects in g_1^d at low x and Q^2 absent ?


Flavour separation of helicity distributions (@ LO)



$$Q^2 = 3 \text{ (GeV/c)}^2$$

Fragmentation functions!


Is the polarised sea flavour symmetric? $\Delta\bar{u} - \Delta\bar{d} = ?$



- COMPASS: $\int_{0.004}^{0.3} (\Delta\bar{u} - \Delta\bar{d}) = 0.052 \pm 0.035 \pm 0.013 @ Q^2 = 3 \text{ GeV}^2$.
- HERMES: $\int_{0.023}^{0.6} (\Delta\bar{u} - \Delta\bar{d}) = 0.048 \pm 0.057 \pm 0.028 @ Q^2 = 2.5 \text{ GeV}^2$.
- Presently accessible only in SIDIS.
- Unpolarised sea: $\int (\bar{u} - \bar{d}) = -0.118 \pm 0.012$

W^\pm production in $\vec{p}\vec{p}$ scattering @ RHIC


$\vec{p}\vec{p} \rightarrow W^{+(-)} \rightarrow l^{+(-)}\nu X$ run 9 @ $\sqrt{s} = 500$ GeV, preliminary



- $\sigma(W)$ agrees reasonably with theory
- Hard scale set by the lepton p_T
- No FF uncertainties

LO interpretation:

- $A_L^{W^-} = \frac{1}{2} \left(\frac{\Delta\bar{u}}{\bar{u}} - \frac{\Delta d}{d} \right) = 0.18 \pm 0.19^{+0.04}_{-0.03}$ (STAR)
- $A_L^{W^+} = \frac{1}{2} \left(\frac{\Delta\bar{d}}{\bar{d}} - \frac{\Delta u}{u} \right) = -0.33 \pm 0.10 \pm 0.04$ (STAR)
 $= -0.83 \pm 0.31$ (PHENIX)
- $x(\Delta\bar{u} - \Delta\bar{d}) > 0$?





After J. Balewski, DIS2010

Global parton analyses


Example: DSSV (D. de Florian, R. Sassot, M. Stratmann, V. Vogelsang),
cf. Phys. Rev. D **80** (2009) 034030.

$$R_q = \frac{\Delta u + \Delta \bar{u}}{u + \bar{u}}$$




- Δg poorly known but small
- ... may still contribute to nucleon spin
- has a node at $x \sim 0.1$?
- $\Delta s > 0$ from SIDIS
but < 0 required by incl. data (+ SU_3)
- ... thus node in Δs fits.

Δg from the polarised pp scattering @ RHIC




PHENIX
(high p_T π^0 production)



STAR
(mid-rapidity inclusive jets)

- Confirmed the electroproduction results
- Included in the DSSV2009.


Δg from the PGF in electroproduction




if $q, \bar{q} \equiv c, \bar{c}$ then $A_{meas} \sim \langle a_{pgf} \rangle \left\langle \frac{\Delta g}{g} \right\rangle + A_{bckg}$ AROMA
 if $q, \bar{q} \equiv \text{light quarks}$ then more MC info needed LEPTO/PYTHIA.

To suppress background in $c\bar{c}$:

- only one charm meson candidate
- quasi-real photons
- A weighting method used to optimise the $\Delta g(x)$ extraction
- Combinatorial background significantly reduced for the $D^* \rightarrow D^0 + \pi_s \rightarrow K + \pi + \pi_s$.




Δg from the PGF in electroproduction, ...cont'd



K.Kurek, COMPASS, DIS2010

Δg from the PGF in electroproduction, ...cont'd



DSSV, arXiv:0904.3821

Nucleon spin decomposition – “22 years later”

$$\frac{1}{2} = \frac{1}{2}\Delta\Sigma + \Delta G + L$$

Are we approaching the solution of the “proton spin puzzle”?

- Restoration of $\Delta\Sigma=0.6$ via the axial anomaly improbable.

As a consequence of the “axial anomaly” the measured quantity is:

$$a_0(Q^2) = \Delta\Sigma^{AB} - \left(\frac{3\alpha_s}{2\pi}\right)\Delta G(Q^2)$$


where COMPASS @ 3 GeV² gives: $a_0 = 0.35 \pm 0.03 \pm 0.05$

and the “spin crisis” can be solved ($\Delta\Sigma \sim 0.6$) if $\Delta G \sim 2.2$ (and $L \sim -2$) at $Q^2 = 3 \text{ GeV}^2$.


- Global, consistent NLO analysis of ΔG needed.
- Independent measurement of L necessary (\implies DVCS, lattice QCD?).
- All candidates are contributing about equally to the nucleon spin?

Partonic structure of the nucleon; distribution functions


Three **twist-two** quark distributions in QCD (after integrating over the quark intrinsic k_t)



Quark momentum DF;
well known (unpolarised DIS $\rightarrow F_{1,2}(x)$).



Difference in DF of quarks with spin parallel or antiparallel to the nucleon's spin;
known (polarised DIS $\rightarrow g_1(x)$).



Difference in DF of quarks with spin parallel or antiparallel to the nucleon's spin in the transversely polarised nucleon;
unknown (polarised DIS $\rightarrow h_1(x)$).

Nonrelativistically: $\Delta_T q(x) \equiv \Delta q(x)$. **OBS.!** $\Delta_T q(x)$ are **C-odd and chiral-odd**; may only be measured with another chiral-odd partner, e.g. fragmentation function.

If the k_t taken into account \implies 8 TMD distr.; one, f_{1T}^\perp accessible through "Sivers asymmetry".

Measurements on a transversely polarised target

Properties of $\Delta_T q(x)$:

- is chiral-odd \implies hadron(s) in final state needed to be observed
- simple QCD evolution since no gluons involved
- related to GPD
- sum rule for transverse spin
- first moment gives “tensor charge” (now being studied on the lattice)

Transversity measured e.g. via the Collins asymmetry (asymmetry in the distribution of hadrons):

$$N_h^\pm(\phi_c) = N_h^0 [1 \pm p_T D_{NN} A_{Coll} \sin \phi_c]$$

which in turn gives at LO:




$$A_{Coll} \sim \frac{\sum_q e_q^2 \cdot \Delta_T q \cdot \Delta_T^0 D_q^h}{\sum_q e_q^2 \cdot q \cdot D_q^h}$$

But **transverse fragmentation functions $\Delta_T^0 D_q^h$ needed to extract $\Delta_T q(x)$** from the Collins asymmetry! Recently those FF measured by BELLE.

Properties of the Sivers process: it is related to L_q in the proton. **Fundamental !**

Results for the transverse asymmetries

COMPASS 2007 proton data



- Collins 1-h asymmetries for proton large at $x \gtrsim 0.1$, consistent with HERMES
- 2-h asymmetry for proton large in the valence region; HERMES sees less.
- Sivers 1-h asymmetries for proton small, contrary to HERMES; needs to be

cleared.

- COMPASS deuteron data: both Collins and Sivers asymmetries very small. These data + Hermes + Belle: $\Rightarrow \Delta_{Tu} + \Delta_{Td} \sim 0$
- First Δ_{Tq} global analyses performed. (cf. Anselmino et al., arXiv:0807.0173)

Outlook

Goals:

- Determination of ΔG .
- Measurements of GPD via DVCS $\longrightarrow L_z$.

Experimental prospects:

- Short term: more lepton data from COMPASS and JLAB
more hadron data from RHIC (500 GeV!)
- 5-10 years: COMPASS DVCS and DY programme
RHIC upgrade
JLAB upgrade 12 GeV.
- ??? years: electron-ion colliders: eRHIC, ELIC, ENC, LHeC

EINSTEIN SIMPLIFIED

