

Search for Heavy (pseudo)Higgs boson A/H produced in association with a top-antitop quark pair leading to the final state with four top quarks in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector

Thursday 17 March 2022 11:40 (20 minutes)

Four top-quark production, a rare process in the Standard Model (SM) with a cross-section around 12 fb, is one of the heaviest final states produced at the LHC, and it is naturally sensitive to physics beyond the Standard Model (BSM). A data excess is observed with twice of the expectation. A follow-up analysis is the search for Heavy (pseudo)Higgs boson A/H produced in association with a top-antitop quark pair leading to the final state with four top quarks. The data analyzed correspond to an integrated luminosity of 139 fb⁻¹. In this talk, the four top-quark decay final states containing either a pair of same-sign leptons or multi-lepton (SSML) are considered. To enhance the search sensitivity, a mass-parameterized BDT is introduced to discriminate the BSM signal against the irreducible SM four-top and other dominant SM backgrounds. Expected upper bounds on the production cross-section of A/H are derived in the mass range from 400 GeV to 1000 GeV.

Career stage

Graduate student

Author: TSAI, Meng-Ju (University of Michigan)

Presenter: TSAI, Meng-Ju (University of Michigan)

Session Classification: BSM I