

## The Stability and Electronic Structure of Magnesium Hydride and Magnesium Deuteride Under High Pressure

The metal polyhydrides have attracted considerable attention because they can evolve into a superconductor with a high value of critical temperature ( $T_c$ ) under pressure. In this research, we have investigated structures of  $MgH_2$ ,  $MgH_3$ , and structures that substitution of deuterium instead of hydrogen under pressure 0-300 GPa to determine the stability of structure under high pressure so that it can be used as a fundamental model for future critical temperature calculations. The calculations are performed by using density functional theory (DFT) based code Quantum Espresso in this work. Generalized gradient approximations (GGA) of Perdew-Burke-Ernzerhof (PBE) have been adopted for exchange-correlation potential. The plane-wave energy cutoff is 80 Ry and the set of K points mesh is  $12 \times 12 \times 12$  for all structures. The band structures reveal the metallic character of the compound. The calculation of energy band structures for  $MgH_2$  and  $MgD_2$  are not different as the same as  $MgH_3$  and  $MgD_3$ . We found that the convex hull of Mg and H have thermodynamically stable at some pressure and the results of phonon calculations confirm that the structure is dynamically stable.

**Primary author:** Mr BOONCHOT, Chayaphon

**Co-authors:** Dr TSUPPAYAKORN-AEK, Prutthipong; Dr BOVORNRATANARAKS, Thiti; Dr PINSOOK, Udomsilp

**Presenter:** Mr BOONCHOT, Chayaphon

**Track Classification:** Condensed Matter Physics