



Workshop

KIAS

Introduction

Section 2

Section 3

Section 4

Conclusions

# Probing $E_6$ Motivated $U(1)$ Theories at the J-PARC KOTO

by F.C.C., P. Ko, S. Baek, C. Yu, Y. Omura

Korea Institute of Advanced Study  
"Higgs as a Probe of New Physics", Osaka University, Japan

March, 2021



Workshop

KIAS

Introduction

Section 2

Section 3

Section 4

Conclusions

- 1 Introduction
- 2 KOTO Searches and  $U(1)_X$  Theories
- 3  $E_6$  Motivated Leptophobic  $U(1)_X$
- 4  $U(1)_X$  Phenomenology
- 5 Conclusions



# Introduction

Workshop

KIAS

Introduction

Section 2

Section 3

Section 4

Conclusions

The recent results provided by the J-PARC KOTO experiment could not improve the upper limit on the  $K_L \rightarrow \pi \bar{\nu} \nu$  branching fraction, given by  $3.0 \cdot 10^{-9}$  at 90% C.L.. Compared to the SM prediction  $\text{Br}(K_L \rightarrow \pi^0 \bar{\nu} \nu) = 0.34(6) \times 10^{-10}$ , it allows a noticeable room for NP which can be investigated from:

$$\begin{aligned} K_L &\rightarrow \varphi_R \varphi_I, & \text{followed by} & \quad \varphi_R \rightarrow \varphi_I + \pi^0; \\ K_L &\rightarrow \bar{\nu}_i \nu_j, & \text{followed by} & \quad \nu_j \rightarrow \nu_i + \pi^0; \\ K_L &\rightarrow \pi^0 Z_H, & \text{followed by} & \quad Z_H \rightarrow \nu + \bar{\nu}. \end{aligned} \quad (1)$$



# KOTO Searches and $U(1)_X$ Theories

Workshop

KIAS

Introduction

Section 2

Section 3

Section 4

Conclusions

$SM \otimes U(1)_X$ : We consider two possible operators for  $\Delta S = 1$   $s \rightarrow d$  transition:

$$O_V = [C_V^L(\overline{s_L}\gamma_\mu d_L) + C_V^R(\overline{s_R}\gamma_\mu d_R)] \times J_X^\mu, \quad (2)$$

$$O_S = [C_S^L(\overline{s_R}d_L) + C_S^R(\overline{s_L}d_R)] \times J_X, \quad (3)$$

where  $J_X^\mu$  and  $J_X$  denote the missing mass current. In order to generate  $O_V$  type operators from NP, the SM fermions must also be chiral under the new  $U(1)_H$  gauge symmetry. Accordingly, there must be at least two important issues to be addressed during the model building step:

- (i) It should be anomaly free, what in general can be achieved by the introduction of new chiral fermions;
- (ii) It should admit renormalizable Yukawa couplings to all the SM fermions. It will commonly require the introduction of additional Higgs doublets charged under  $U(1)_H$ .



# $E_6$ Motivated Leptophobic $U(1)_X$

Workshop

KIAS

Introduction

Section 2

Section 3

Section 4

Conclusions

We seek for a general charge assignment of a particle content satisfying the following conditions:

- 1 The representations define a  $E_6 SM \otimes U(1)_H$  subgroup for at least one of the collection of models we construct. It will correspond to extend the SM content with the  $SU(2)_L$  neutral lepton singlets  $n_R$  and  $N_L$ , a pair of d-type quark singlets  $D_{L(R)}$ , and the lepton doublets  $L_{L(R)}$ ;
- 2 The  $h$ -charges constrain the Yukawa interactions between the SM fermions to those of a type-II 2HDM;
- 3  $U(1)_H$  is leptophobic over SM leptons.



# $E_6$ Motivated Leptophobic $U(1)_X$

Workshop

KIAS

Introduction

Section 2

Section 3

Section 4

Conclusions

	$q_L^i$	$u_R^i$	$d_R^i$	$l_L^i$	$e_R^i$	$n_R^i$	$H_1$
$SU(3)_c$	3	3	3	1	1	1	1
$SU(2)$	2	1	1	2	1	1	2
$U(1)_Y$	1/6	2/3	-1/3	-1/2	-1	0	1/2
$U(1)_H$	$b(1+a)$	$-2b(1+a)$	$b(1+a)$	0	0	$b\left(\pm\sqrt{\frac{a(9+14a)}{2}} - 3(1+a)\right)$	0

TABLE I: General charge assignments of the SM fermions under the SM gauge group and the Leptophobic  $U(1)_H$   $E_6$  subgroup.

	$H_2$	$D_L^i$	$D_R^i$	$L_L^i$	$L_R^i$	$N_L^i$	$\Phi_k$	$\varphi$
$SU(3)_c$	1	3	3	1	1	1	1	1
$SU(2)$	2	1	1	2	2	1	1	1
$U(1)_Y$	1/2	-1/3	-1/3	-1/2	-1/2	0	0	0
$U(1)_H$	$-3b(1+a)$	$-b(2+3a)$	$b$	$ab$	$b(3+4a)$	$b\left(\pm\sqrt{\frac{a(9+14a)}{2}} + 3(1+a)\right)$	$X_k$	$X_\varphi$

TABLE II: Charge of the new particle content under the SM gauge group and the Leptophobic  $U(1)_H$   $E_6$  subgroup.



# Complete Example

Workshop

KIAS

Introduction

Section 2

Section 3

Section 4

Conclusions

$a = -6$ :

$$\begin{aligned} \mathcal{L} \supset & Y_\nu \bar{l}_L \tilde{H}_1 n_R + M_n \bar{n}_R^c n_R + Y_{2N} \Phi_2 \bar{n}_R N_L \\ & + Y_L \Phi^\dagger \bar{L}_L L_R + Y_D \Phi^\dagger \bar{D}_L D_R + Y_{l\chi} \varphi_\chi \bar{l}_L L_R + Y_{d\chi} \varphi_\chi \bar{D}_L d_R + \text{h.c.}, \end{aligned} \quad (4)$$

Notice that  $c_\beta = v_1/v$  is supposed to modulate neutrino masses.

$$\begin{aligned} V(H_i, \Phi, \varphi) \supset & m_\varphi^2 |\varphi|^2 + \lambda_\varphi |\varphi|^4 + \lambda_{i\varphi} |\varphi|^2 H_i^\dagger H_i + \lambda_{\Phi\varphi} |\varphi|^2 |\Phi|^2 \\ & + \left\{ (\lambda_{12\varphi} H_1^\dagger H_2 + \sqrt{2} \mu_{\Phi\varphi} \Phi^\dagger) \varphi^2 + \text{H.c.} \right\} \end{aligned}$$

$\varphi_I$ : Light DM candidate

$$m_I^2 = m_R^2 - 2 (\lambda_{12\varphi} v^2 c_\beta s_\beta + 2 \mu_{\Phi\varphi} v_\Phi), \quad (5)$$

# KOTO Searches from Dark Fields

Workshop

KIAS


Introduction

Section 2

Section 3

Section 4

Conclusions



- Rare processes are typically related to  $\Delta F = 1, 2$ .
- Universal theories: FCNC at tree-level only from mixing between SM and NP;
- For DM theories, i.e. if NP sector is dark ( $Z_2$  odd), the physics of portals will emerge at the loop-level: Need of generalized formulas for penguins, for both vector and pseudo-scalar mediators;

$$i\Gamma_A = g^2 \frac{V_{us}^* V_{ud}}{16\pi^2} A \{ \alpha_R [\bar{s}_L d_R] + \alpha_L [\bar{s}_R d_L] \} \quad (6)$$

$$i\Gamma_G = ig^2 \frac{V_{us}^* V_{ud}}{16\pi^2} z_L^G G_\mu [\bar{s}_L \gamma^\mu d_L]. \quad (7)$$





# $U(1)_H$ Phenomenology

Workshop

KIAS


Introduction

Section 2


Section 3

Section 4


Conclusions




(Ia)




(Ib)




(IIa)




(IIIb)




(IIIa)



(IVa)



(IVb)



(VI)

Feynman diagrams for  $K_L \rightarrow \phi_I \phi_R$ .



# $U(1)_H$ Phenomenology

Workshop

KIAS

Introduction

Section 2



Section 3

Section 4

Conclusions

- i Pseudo-scalar  $A$  only, where  $Z_H$  effects are absent and  $s_\theta \rightarrow 0$ . This is close to test 2HDM models;
- ii  $Z, Z_H$  only and a decoupled pseudo-scalar. This is close to test  $U(1)_X$  theories;
- iii All off-shell  $A, Z, Z_H$  contributing. This scenario tests  $2HDM_X$ , and includes massive neutrinos as the source of missing mass;
- iv Light  $Z_H$  resonance mediating fermion production ( $\nu_i$ ).

# $U(1)_H$ Phenomenology



$K_L \rightarrow \varphi_R \varphi_I$  at the EW scale, mediated by (a) pseudo-scalar and (b)  $G - Z, Z_H$ .

- At the EW scale, the theory would rely on the presence of a light (MeV) scalar among its spectra, which must regulate  $\varphi_I$  abundance;
- DM annihilation from light scalar, s-channel: p-wave, safe from CMB bounds;
- The light scalar would enter in the KOTO physics at the three-body level;



# $U(1)_H$ Phenomenology

Workshop

KIAS



Introduction

Section 2

Section 3

Section 4

Conclusions



$K_L \rightarrow \varphi_R \varphi_I$  and  $K_L \rightarrow \nu_i \nu_j$ . The red band is affected by the mass-mixing suppression in leptophobic theories.

- $\varphi_I$  abundance regulated by  $\varphi_I \varphi_R \rightarrow Z_H^* \rightarrow SM.$ ;
- $\varphi_R$  decayed away at time of recombination: safe from CMB bounds;
- The  $\nu_j \rightarrow \nu_i \pi$  requires the hierarchy  $(M_{\nu_i}, M_{\nu_j}, M_{\nu_k}) = (0.1\text{eV}, \text{MeV}, M_{\nu_k})$ , which can be realized in our setup;



# Conclusions

Workshop

KIAS

Introduction

Section 2

Section 3

Section 4

Conclusions

- We provide a class of theories of enlarged dark sector;
- We provide generalized formulas of loop-corrected  $\Delta F = 1$  vertices;
- We probe the physics of vector and pseudo-scalar portals for a particular DM model at the J-PARC KOTO.