

Probing EWPT in 2HDM with Future Lepton Colliders

Wei Su

University of Adelaide

arXiv: [2011.04540](https://arxiv.org/abs/2011.04540) WS, A G. Williams, M. Zhang



THE UNIVERSITY
of ADELAIDE

Outline

 2HDM and Phase Transition



 Higgs/Z-pole Precision Measurements

 Results: cases and general scan

 Conclusion

Electroweak Phase Transition

baryon asymmetry of the Universe (BAU)



SM: Cross-over around $T=100$ GeV

BSM: bubble formation \longrightarrow asymmetry

2HDM: Brief Introduction

- Two Higgs Doublet Model

$$\Phi_i = \begin{pmatrix} \phi_i^+ \\ (v_i + \phi_i^0 + iG_i)/\sqrt{2} \end{pmatrix}$$

$$v_u^2 + v_d^2 = v^2 = (246\text{GeV})^2$$

$$\tan \beta = v_u/v_d$$

	ϕ_1	ϕ_2
Type I	u,d,l	
Type II	u	d,l
lepton-specific	u,d	l
flipped	u,l	d

$$\begin{pmatrix} H^0 \\ h^0 \end{pmatrix} = \begin{pmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} \phi_1^0 \\ \phi_2^0 \end{pmatrix},$$

$$A = -G_1 \sin \beta + G_2 \cos \beta$$

$$H^\pm = -\phi_1^\pm \sin \beta + \phi_2^\pm \cos \beta$$

- Parameters (CP-conserving, Flavor Limit, Z_2 Symmetry)

$$m_{11}^2, m_{22}^2, \lambda_1, \lambda_2, \lambda_3, \lambda_4, \lambda_5$$



$$v, \tan \beta, \alpha, m_h, m_H, m_A, m_{H^\pm}$$


Soft Z_2 symmetry breaking: m_{12}^2

246 GeV

125. GeV


2HDM: precision

Model	κ_V	κ_u	κ_d	κ_ℓ
2HDM-I	$\sin(\beta - \alpha)$	$\cos \alpha / \sin \beta$	$\cos \alpha / \sin \beta$	$\cos \alpha / \sin \beta$
2HDM-II	$\sin(\beta - \alpha)$	$\cos \alpha / \sin \beta$	$-\sin \alpha / \cos \beta$	$-\sin \alpha / \cos \beta$
2HDM-L	$\sin(\beta - \alpha)$	$\cos \alpha / \sin \beta$	$\cos \alpha / \sin \beta$	$-\sin \alpha / \cos \beta$
2HDM-F	$\sin(\beta - \alpha)$	$\cos \alpha / \sin \beta$	$-\sin \alpha / \cos \beta$	$\cos \alpha / \sin \beta$



$$\lambda v^2 \equiv m_\Phi^2 - m_{12}^2/s_\beta c_\beta$$



$$-125^2 \text{GeV}^2 < \lambda v^2 < 600^2 \text{GeV}^2$$



2HDM: precision

[1808.02037](#) N. Chen, T. Han, S. Su, WS, Y. Wu


Type-II, CEPC



2HDM: precision


[1808.02037](#) N. Chen, T. Han, S. Su, WS, Y. Wu

Type-II, CEPC





[arXiv:1405.5537](#) G. C. Dorsch, S. Huber, K. Mimasu, J. M. No


Discovering a second Higgs doublet through $A_0 \rightarrow H_0 Z$.



2HDM: LHC direct search



Results: Case-1



Type-II
fixed mass splitting 200 GeV

$m_H < 710$ GeV
 $\tan\beta \in (1.8, 10)$

Vacuum uplifting:



[arXiv:1705.09186](https://arxiv.org/abs/1705.09186)

[G. C. Dorsch](#), [S. Huber](#), [K. Mimasu](#), [J. M. No](#)

$$\Delta\mathcal{F}_0 = \frac{1}{64\pi^2} \left[(m_h^2 - 2M^2)^2 \left(\frac{3}{2} + \frac{1}{2} \log \left[\frac{4m_A m_H m_{H^\pm}^2}{(m_h^2 - 2M^2)^2} \right] \right) + \frac{1}{2} (m_A^4 + m_H^4 + 2m_{H^\pm}^4) + (m_h^2 - 2M^2) (m_A^2 + m_H^2 + 2m_{H^\pm}^2) \right]$$

Results: Case-2


$$m_A = m_{H^\pm} \tan \beta = 3$$




Too large masses or mass splitting can not generate SFOEWPT

Results: Case-2/3

$$m_A = m_{H^\pm} \tan \beta = 3$$





$$m_H = 700 \text{ GeV}$$





Too large masses or mass splitting can not generate SFOEWPT

Results: Type-II





Future






[1812.01633](#)
 F. Kling, H. Li, etc

Results: Type-I





Conclusion





Thanks!

PT vs. vacuum uplifting

$$\xi_c \equiv \frac{v_c}{T_c}$$




Future



[1812.01633](#)
 F. Kling, H. Li, etc

2HDM:Theoretical constraints



$$\cos(\beta - \alpha) = 0$$

$$m_\Phi \equiv m_H = m_A = m_{H^\pm}$$

$$\lambda v^2 \equiv m_\Phi^2 - m_{12}^2 / s_\beta c_\beta$$

$$-125^2 \text{ GeV}^2 < \lambda v^2 < 600^2 \text{ GeV}^2$$

$$\lambda \in (-0.26, 5.95)$$

$$\lambda_4 = \lambda_5 = \lambda_3 - 0.258 = -\lambda$$

Results: Case-2/3

High T approximation:

$$V(\phi_h, T) \approx (DT^2 - \mu^2)\phi_h^2 - ET\phi_h^3 + \frac{\tilde{\lambda}}{4}\phi_h^4$$

$$D = \frac{1}{24} \left[6\frac{m_W^2}{v^2} + 3\frac{m_Z^2}{v^2} + \frac{m_h^2}{v^2} + 6\frac{m_t^2}{v^2} + \frac{m_H^2 - M^2}{v^2} + \frac{m_A^2 - M^2}{v^2} + 2\frac{m_{H^\pm}^2 - M^2}{v^2} \right]$$

$$E = \frac{1}{12\pi} \left[6\frac{m_W^3}{v^3} + 3\frac{m_Z^3}{v^3} + \frac{m_h^3}{v^3} \right] + E_{(H/A/H^\pm)}$$

$$E_{(\alpha)} \approx \begin{cases} \frac{1}{12\pi} \lambda_\alpha^{3/2} = \frac{1}{12\pi} \frac{m_\alpha^3}{v^3}, & M^2 \ll \lambda_\alpha \phi_h^2 \\ 0, & M^2 \gg \lambda_\alpha \phi_h^2 \end{cases}$$

$$\lambda_{A/H^\pm} v^2 = (\Delta m)^2 + 2m_H \Delta m$$

Vacuum uplifting:

$$\Delta \mathcal{F}_0 = \frac{1}{64\pi^2} \left[(m_h^2 - 2M^2)^2 \left(\frac{3}{2} + \frac{1}{2} \log \left[\frac{4m_A m_H m_{H^\pm}^2}{(m_h^2 - 2M^2)^2} \right] \right) \right. \\ \left. + \frac{1}{2} (m_A^4 + m_H^4 + 2m_{H^\pm}^4) + (m_h^2 - 2M^2) (m_A^2 + m_H^2 + 2m_{H^\pm}^2) \right]$$

Too large masses or mass splitting can not generate SFOEWPT