

Dark Photon from Scalar Boson Decays at FASER

Takashi Shimomura
(Miyazaki U.)

based on

“Dark Photon from Light Scalar Boson Decays at FASER”
JHEP03 (2021), arXiv:2008.12765

in collaboration with

Takeshi Araki	(Ohu U.)
Kento Asai	(Tokyo U.)
Hidetoshi Otono	(Kyushu U.)
Yosuke Takubo	(KEK)

Introduction

- ▶ Non-observation of Dark Matter/BSM leads to the idea of “**dark sector**”.



- ▶ No direct interactions between the SM and dark particles.
- ▶ Messenger particle (portal) connects two sectors.

Portals

Vector portal, Scalar portal, Heavy neutral lepton portal, Axion portal
dark photon dark higgs


Dark Photon

- ▶ Gauge boson of a new $U(1)_{\text{dark}}$ symmetry.
- ▶ Mixing with $U(1)_Y$ or $U(1)_{\text{EM}}$ gauge boson

$$\mathcal{L} = -\frac{\epsilon}{2} F^{\mu\nu} F'_{\mu\nu}$$


ϵ : gauge kinetic mixing

- ▶ Dark Photon (DP, A') can interact with the SM particles **via ϵ** .



$$\mathcal{L}_{\text{int}} \supset \epsilon e A'_\mu J_{\text{EM}}^\mu$$

(couple to the EM current)



Dark photon can be long-lived

$$\epsilon \sim 10^{-7}$$



$$c\tau \sim O(100) \text{ m}$$

Origin of DP mass

- ▶ One of the dynamical origins of the mass is **spontaneous breaking**.
 - ▶ A new scalar boson should be introduced to give a mass.

Dark Higgs kinetic term

$$\mathcal{L} \supset g'^2 \phi^\dagger \phi A'_\mu A'^\mu \xrightarrow[\langle \phi \rangle = v_\phi / \sqrt{2}]{\text{U(1)}_{\text{dark br.}}} \frac{1}{2} m_{A'}^2 A'_\mu A'^\mu + g' m_{A'} \phi A'_\mu A'^\mu$$

Direct consequence of the mass generation.

Origin of DP mass

- ▶ One of the dynamical origins of the mass is **spontaneous breaking**.
 - ▶ A new scalar boson should be introduced to give a mass.

Dark Higgs kinetic term

$$\mathcal{L} \supset g'^2 \phi^\dagger \phi A'_\mu A'^\mu \xrightarrow[\langle \phi \rangle = v_\phi / \sqrt{2}]{U(1)_{\text{dark br.}}} \frac{1}{2} m_{A'}^2 A'_\mu A'^\mu + g' m_{A'} \phi A'_\mu A'^\mu$$

Direct consequence of the mass generation.

- ▶ Dark Higgs decays into $A'A'$
 - ▶ Controlled by g' , not ε .
 - ▶ **Enhanced when $m_{A'} \ll m_\phi$.**



$$\Gamma(\phi \rightarrow A'A') = \frac{g'^2}{8\pi} \frac{m_{A'}^2}{m_\phi} \beta_\phi(A') \left[2 + \frac{m_\phi^4}{4m_{A'}^4} \left(1 - \frac{2m_{A'}^2}{m_\phi^2} \right)^2 \right]$$

$$\text{where } \beta_\phi(A') = \sqrt{1 - 4m_{A'}^2/m_\phi^2}$$

FASER experiment

Feng, Galon, Kling, Trojanowski, PRD97 (2018)
 FASER collaboration, arXiv:1708.09389

- ▶ ForwArd Search ExpeRiment (FASER) at LHC, starting from 2022.
- ▶ Detector is placed 480m downstream from the ATLAS interaction point.
- ▶ Search for **dark photon**, **dark Higgs**, Axion-like particle, etc.




New light and weakly int. particles preferentially go in forward direction

	L_{\min} (m)	L_{\max} (m)	R (m)	\mathcal{L} (ab^{-1})
FASER	478.5	480	0.1	0.15
FASER 2	475	480	1.0	3.0

DP search at FASER


- Production of dark photon
 - meson decays : $\pi^0 \rightarrow \gamma A'$, $\eta \rightarrow \gamma A'$
 - proton bremsstr. : $pp \rightarrow p X A'$
 - QCD processes : $q\bar{q} \rightarrow g A'$, $qg \rightarrow q A'$
- Detection
 - $A' \rightarrow e\bar{e}, \mu\bar{\mu}, \text{hadrons}$



DP search at FASER

- Production of dark photon
 - meson decays : $\pi^0 \rightarrow \gamma A'$, $\eta \rightarrow \gamma A'$
 - proton brems. : $pp \rightarrow p X A'$
 - QCD processes : $q\bar{q} \rightarrow g A'$, $qg \rightarrow q A'$
- Detection
 - $A' \rightarrow e\bar{e}, \mu\bar{\mu}, \text{hadrons}$

FASER/FASER 2 can explore parameter region between collider and beam dump exp.



Dark Higgs Production

Feng, Galon, Kling, Trojanowski, PRD97 (2018)


- Dark Higgs is produced from meson decays through the ϕ -Higgs mixing.

meson decays :

$$B \rightarrow X_s \phi$$



$$K \rightarrow \pi \phi$$

$$\eta' \rightarrow \eta \phi$$



more than 100-1000 DH can be produced

(0.3 GeV, 4×10^{-4})




Dark Higgs Decay

Assuming $m_\phi \gg m_{A'}, m_f$,

$$\Gamma(\phi \rightarrow A'A') \simeq \frac{g'^2}{32\pi} \left(\frac{m_\phi}{m_{A'}} \right)^2 m_\phi$$

$$\Gamma(\phi \rightarrow f\bar{f}) \simeq \frac{\theta^2}{8\pi} \left(\frac{m_f}{v} \right)^2 m_\phi$$



Dark Higgs Decay

Assuming $m_\phi \gg m_{A'}, m_f$,

$$\Gamma(\phi \rightarrow A'A') \simeq \frac{g'^2}{32\pi} \left(\frac{m_\phi}{m_{A'}} \right)^2 m_\phi$$

$$\Gamma(\phi \rightarrow f\bar{f}) \simeq \frac{\theta^2}{8\pi} \left(\frac{m_f}{v} \right)^2 m_\phi$$

- ▶ The ϕ mainly decays into DPs for $g' > \theta$.



New source of DPs

→ smaller ε can be searched

- ▶ The ϕ can be long-lived for $g' \ll \theta$.

Carrier of dark photon

→ larger ε can be searched



Short-lived DH at FASER2



95% C.L. (>3)
projected sensitivity
only from direct prod.

(FASER, PRD99,9, (2019))

$$\pi^0 \rightarrow \gamma A', \eta \rightarrow \gamma A'$$


$$pp \rightarrow p X A'$$

$$q\bar{q} \rightarrow g A', qg \rightarrow q A'$$

NuCal, CHARM,
LSND, etc.

Supernova cooling

KLOE, BaBar, NA48/2, NA64, etc.



Number of A' decays
produced from ϕ

>100

10-99

3-9

Short-lived DH at FASER2



95% C.L. (>3)
projected sensitivity
only from direct prod.

(FASER, PRD99,9, (2019))

$$\pi^0 \rightarrow \gamma A', \eta \rightarrow \gamma A'$$


$$pp \rightarrow p X A'$$

$$q\bar{q} \rightarrow g A', qg \rightarrow q A'$$

NuCal, CHARM,
LSND, etc.

Supernova cooling

KLOE, BaBar, NA48/2, NA64, etc.



Number of A' decays
produced from ϕ

>100


10-99

3-9

Signal of DP production
from ϕ decay

The origin of the DP mass
can be searched.

Long-lived DH at FASER2



95% C.L. (>3)
projected sensitivity
only from direct prod.

(FASER, PRD99,9, (2019))

$$\pi^0 \rightarrow \gamma A', \eta \rightarrow \gamma A'$$


$$pp \rightarrow p X A'$$

$$q\bar{q} \rightarrow g A', qg \rightarrow q A'$$

NuCal, CHARM,
LSND, etc.

Supernova cooling

KLOE, BaBar, NA48/2, NA64, etc.



Signal of DP production
from long-lived ϕ decay




Number of A' decays
produced from ϕ

>100

10-99

3-9

Lighter Dark Higgs





Summary

- ▶ We consider dark photon and dark higgs as the origin of DP mass.
 - The dark higgs- $A'-A'$ interaction appears as the consequence of symmetry breaking.
 - The signal can be found in different region from directly produced DPs both in short-lived or long-lived dark Higgs cases.
 - The origin of DP mass will be investigated at FASER(2).


Thank you for your attention !

Back-Up Slides

Long-lived at FASER2



decay probability :



Some DP decay inside the detector even it is long-lived.

Signal Calculation

- ▶ The ϕ production from mesons is only considered.
- ▶ We have calculated the signal of DP decays.
(using data of B meson production provided by Felix Kling, thanks!)

$$\begin{aligned}
 N &= \mathcal{L} \int d\mathbf{p}_{A'} \frac{d\sigma_{pp \rightarrow A'X}}{dp_{A'} d\theta_{A'}} \mathcal{P}_{A'}^{\text{det}}(\mathbf{p}_{A'}, \mathbf{p}_{\phi}) \\
 &= \mathcal{L} \sum_{i:\text{meson}} \sum_{j=1,2} \int dp_i d\theta_i \int d\mathbf{p}_{A'} \int d\mathbf{p}_{\phi} \frac{d\sigma_{pp \rightarrow iX}}{dp_i d\theta_i} \text{Br}(i \rightarrow \tilde{X}\phi) \text{Br}(\phi \rightarrow A'_1 A'_2) \\
 &\quad \times \mathcal{P}_{A'_j}^{\text{det}}(\mathbf{p}_{A'}, \mathbf{p}_{\phi}) ,
 \end{aligned}$$

with

$$\begin{aligned}
 \text{decay probability : } \mathcal{P}_{A'}^{\text{det}}(\mathbf{p}_{A'}, \mathbf{p}_{\phi}) &= \frac{1}{\bar{d}_{\phi} \cos \theta_{\phi}} \int_{z_{\phi,\min}}^{z_{\phi,\max}} dz_{\phi} e^{-\frac{z_{\phi}}{\bar{d}_{\phi} \cos \theta_{\phi}}} \frac{1}{\bar{d}_{A'} \cos \theta_{A'}} \int_{z_{A',\min}}^{L_{\max}} dz_{A'} e^{-\frac{z_{A'} - z_{\phi}}{\bar{d}_{A'} \cos \theta_{A'}}} \\
 &\quad \times \Theta(R - r_{A',R}) \Theta(R - r_{A',F}),
 \end{aligned}$$

- ▶ Cut $p_{A'} > 100$ GeV is imposed.

Beam Dump Exp.

Special thanks Felix and Sebastian

For $m_\phi = 2$ GeV, $\theta=10^{-4}$

▶ CHARM experiment

- proton beam energy : 400 GeV

$$N_{\text{pot}} = 2 \times 10^{18} \longrightarrow N_{\phi \text{ total}} = 3.0 \times 10^4 \quad (N_B = 6.4 \times 10^{11})$$

Alekhin et al, 1504.04855

- The detection probability

$$P \sim 10^{-4} \text{ for } 0.068 < \theta_{\text{det}} < 0.0126$$

Dobrich et al, 1512.03069

$$\longrightarrow N_{A'} = 6.0 \times 10^4 \times 10^{-4} = 6$$

▶ NuCal experiment

- proton beam energy : 70 GeV

$$N_{\text{pot}} = 1.7 \times 10^{18} \longrightarrow N_{\phi \text{ total}} = O(10) \quad (\text{Bremsstrahlung})$$

Batell et al, 2008.08108

- The detection probability

$$P \sim 10^{-1} \text{ for } \theta_{\text{det}} < 0.015$$

$$\longrightarrow N_{A'} = O(1)$$

might not be significant, but need more detailed analyses