



UNIVERSIDAD  
DE COLIMA



# Predictions for the Leptonic Dirac CP-Violating Phase


Alexander J. Stuart

May 26, 2021

Pheno 2021

Based on: L.A. Delgadillo, L.L. Everett, R. Ramos and AS,  
Phys.Rev. D97 (2018) no.9, 095001 [arXiv:1801.06377] and  
L.L Everett, R. Ramos, A.B. Rock, and AS [arXiv:1912.10139]

# What We Measure



## Quark Mixing

## Lepton Mixing

$$U_{CKM} = R_1(\theta_{23}^{CKM})R_2(\theta_{13}^{CKM}, \delta_{CKM})R_3(\theta_{12}^{CKM}) \quad U_{MNSP} = R_1(\theta_{23})R_2(\theta_{13}, \delta_{CP})R_3(\theta_{12})P$$

$$\theta_{13}^{CKM} = 0.2^\circ \pm 0.1^\circ$$

$$\theta_{13}^{MNSP} = (8.61^\circ)_{-0.13}^{+0.13}$$

NuFIT 4.1 (2019):  
1811.05487

$$\theta_{23}^{CKM} = 2.4^\circ \pm 0.1^\circ$$

$$\theta_{23}^{MNSP} = (48.3^\circ)_{-1.9}^{+1.1}$$

$$\theta_{12}^{CKM} = 13.0^\circ \pm 0.1^\circ$$

$$\theta_{12}^{MNSP} = (33.82^\circ)_{-0.76}^{+0.78}$$

$$\delta_{CKM} = 60^\circ \pm 14^\circ$$

$$\delta_{CP} = (222^\circ)_{-28}^{+38}$$

Quarks look like deviations from unity.

What about the leptons?



# Popular Patterns (before 2012)

$$\tilde{U}_\nu = R_{23}(\theta_{23}^\nu) R_{12}(\theta_{12}^\nu) = \begin{pmatrix} \cos \theta_{12}^\nu & \sin \theta_{12}^\nu & 0 \\ -\frac{\sin \theta_{12}^\nu}{\sqrt{2}} & \frac{\cos \theta_{12}^\nu}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ -\frac{\sin \theta_{12}^\nu}{\sqrt{2}} & \frac{\cos \theta_{12}^\nu}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix} \quad \begin{array}{l} \text{(Marzocca, et al. (2013);} \\ \text{Petcov (2014);} \\ \text{Girardi, Petcov, Titov (2015))} \end{array}$$

TriBiMaximal (TBM) Mixing:  $\theta_{12}^\nu \approx 35.26^\circ$  (P. Harrison, D. Perkins, W. Scott (2002); Z. Xing (2002); X. He, A. Zee (2003))

BiMaximal (BM) Mixing:  $\theta_{12}^\nu = 45^\circ$  (F. Vissani (1997); V. Barger, S. Pakvasa, T. Weiler, K. Whisnant, (1998); A. Baltz, A. Goldhaber, M. Goldhaber (1998))

Golden Ratio 1 (GR1) Mixing:  $\theta_{12}^\nu \approx 31.72^\circ$  (A. Datta, F. Ling, P. Ramond (2003); L. Everett, AS (2008))

Golden Ratio 2 (GR2) Mixing:  $\theta_{12}^\nu = 36^\circ$  (W. Rodejohann (2009))

Hexagonal (HG) Mixing:  $\theta_{12}^\nu = 30^\circ$  (C. Albright, A. Dueck and W. Rodejohann (2010); J. E. Kim and M. Seo (2011))

However, they all have a vanishing reactor mixing angle  $\theta_{13}$ . How can we fix this?

# Charged Lepton Corrections

$$U_{\text{MNSP}} = U_e^\dagger U_\nu$$

As on previous slide, assume the neutrino mixing matrix is given as  $U_\nu = R_{23}(\theta_{23}^\nu)R_{12}(\theta_{12}^\nu)$ .

$$R_{23}^\nu = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23}^\nu & s_{23}^\nu \\ 0 & -s_{23}^\nu & c_{23}^\nu \end{pmatrix} \quad R_{12}^\nu = \begin{pmatrix} c_{12}^\nu & s_{12}^\nu & 0 \\ -s_{12}^\nu & c_{12}^\nu & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Postulate a theoretical form for  $U_e$ , *i.e.*, one rotation, two rotations, etc. For simplicity assume  $U_e$  consists of only a 1-2 rotation:

$$U_{12}^e = \begin{pmatrix} c_{12}^e & s_{12}^e e^{-i\delta_{12}^e} & 0 \\ -s_{12}^e e^{i\delta_{12}^e} & c_{12}^e & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad s_{ij}^e = \sin \theta_{ij}^e \quad \text{and} \quad c_{ij}^e = \cos \theta_{ij}^e$$

How does this charged lepton rotation change the initial mixing predictions?

$$U_e = U_{12}^e(\theta_{12}^e, \delta_{12}^e)$$

# The Effect of a 1-2 Rotation

$$U_{\text{MNSP}} \equiv U = U_e^\dagger U_\nu = U_{12}^{e\dagger} R_{23}^\nu R_{12}^\nu$$

$$\begin{aligned} U_{e1} &= c_{12}^e c_{12}^\nu + c_{23}^\nu e^{-i\delta_{12}^e} s_{12}^e s_{12}^\nu, & U_{e2} &= c_{12}^e s_{12}^\nu - c_{12}^\nu c_{23}^\nu e^{-i\delta_{12}^e} s_{12}^e, \\ U_{e3} &= -e^{-i\delta_{12}^e} s_{12}^e s_{23}^\nu, & U_{\mu1} &= -c_{12}^e c_{23}^\nu s_{12}^\nu + c_{12}^\nu e^{i\delta_{12}^e} s_{12}^e \\ U_{\mu2} &= c_{12}^e c_{12}^\nu c_{23}^\nu + e^{i\delta_{12}^e} s_{12}^e s_{12}^\nu, & U_{\mu3} &= c_{12}^e s_{23}^\nu, \\ U_{\tau1} &= s_{12}^\nu s_{23}^\nu, & U_{\tau2} &= -c_{12}^\nu s_{23}^\nu, \\ U_{\tau3} &= c_{23}^\nu. \end{aligned}$$

Compare this to the PDG parameterization of the MNSP Matrix:

$$U^{\text{PDG}} = \begin{pmatrix} c_{12}c_{13} & c_{13}s_{12} & e^{-i\delta}s_{13} \\ -c_{23}s_{12} - c_{12}e^{i\delta}s_{13}s_{23} & c_{12}c_{23} - e^{i\delta}s_{12}s_{13}s_{23} & c_{13}s_{23} \\ s_{12}s_{23} - c_{12}c_{23}e^{i\delta}s_{13} & -c_{12}s_{23} - c_{23}e^{i\delta}s_{12}s_{13} & c_{13}c_{23} \end{pmatrix} P_{\text{Maj}}$$

By equating both parameterizations, it should become clear that it is possible to express the PDG parameters in terms of the model parameters....

# Enter Sum Rules

Specifically, one can find a relationship between the Dirac CP-violating phase  $\delta$ , the experimentally measured PDG matrix, and the model parameters, i.e.,

$$\cos \delta = \frac{(1/t_{23} + s_{13}^2 t_{23})(s_{12}^\nu)^2 - (s_{12}^2/t_{23} + c_{12}^2 s_{13}^2 t_{23})}{s'_{12} s_{13}}$$

(D. Marzocca, S.T. Petcov, et al. (2011);  
D. Marzocca, S.T. Petcov, et al. (2013);  
P. Ballet, S.F. King, et al. (2014)  
S.T. Petcov (2015);  
I. Girardi, S.T. Petcov, et al. (2015);)

Can be derived easily by taking the ratio (P. Ballet, S.F. King, et al. (2014)):

$$\frac{|U_{\tau 1}|}{|U_{\tau 2}|} = t_{12}^\nu \quad \frac{|U_{\tau 1}^{PDG}|}{|U_{\tau 2}^{PDG}|} = \frac{|s_{12} s_{23} - c_{12} c_{23} s_{13} e^{i\delta}|}{|c_{12} s_{23} + c_{23} s_{12} s_{13} e^{i\delta}|}$$

This sum rule can in principle be applied to reveal how this 1-2 rotation affects the predictions for the leptonic Dirac CP-violating phase. In principle there exists  $9!/2!7!-1=35$  other additional ratios that must hold to guarantee the unitarity of the matrices.

# A Simple set of 4 Sum Rules

The 36 sum rules, upon equating parameters, reduce to only 4 sum rules. The original:

$$\cos \delta = \frac{(1/t_{23} + s_{13}^2 t_{23})(s_{12}^\nu)^2 - (s_{12}^2/t_{23} + c_{12}^2 s_{13}^2 t_{23})}{s_{12}' s_{13}}$$

And 3 others which give the angles of the MNSP matrix in terms of model parameters:


$$s_{13}^2 = (s_{12}^e)^2 (s_{23}^\nu)^2, \quad s_{23}^2 = \frac{(s_{23}^\nu)^2 - (s_{12}^e)^2 (s_{23}^\nu)^2}{1 - (s_{12}^e)^2 (s_{23}^\nu)^2},$$
$$s_{12}^2 = \frac{(c_{12}^\nu)^2 (c_{23}^\nu)^2 (s_{12}^e)^2 + (c_{12}^e)^2 (s_{12}^\nu)^2 - 2c_{12}^e c_{12}^\nu c_{23}^\nu s_{12}^e s_{12}^\nu \cos \delta_{12}^e}{1 - (s_{12}^e)^2 (s_{23}^\nu)^2}$$

The value of  $\cos \delta$  is subject to the constraints in these 3 additional sum rules. How does this value look for the different mixing patterns:

# BM Mixing and a 1-2 Rotation

It is possible to showcase the allowed regions of  $\cos(\delta)$  by using a contour plot:

$$\cos \delta(\theta_{12}^e, \delta_{12}^e)$$



$$\sin^2(\theta_{13}) \equiv s_{13}^2 = (s_{12}^e)^2/2$$

$$s_{23}^2 = \frac{(c_{12}^e)^2/2}{(1 - (s_{12}^e)^2/2)} = \frac{1 - 2s_{13}^2}{2(1 - s_{13}^2)}$$

$$0.4878 \leq s_{23}^2 \leq 0.4904$$

**(First octant!)**

Blue bands and regions between red contours represent regions allowed by  $\sin^2(\theta_{13})$  and  $\sin^2(\theta_{12})$  at  $3\sigma$ , respectively. Notice  $\cos(\delta) \sim -1$  is preferred value.

How about for another popular mixing scenario?

# TBM Mixing and a 1-2 Rotation

$$\cos \delta(\theta_{12}^e, \delta_{12}^e)$$


Blue bands and regions between red contours represent regions allowed by  $\sin^2(\theta_{13})$  and  $\sin^2(\theta_{12})$  at  $3\sigma$ , respectively.

$$\sin^2(\theta_{13}) \equiv s_{13}^2 = (s_{12}^e)^2/2$$

$$s_{23}^2 = \frac{(c_{12}^e)^2/2}{(1 - (s_{12}^e)^2/2)} = \frac{1 - 2s_{13}^2}{2(1 - s_{13}^2)}$$


$$0.4878 \leq s_{23}^2 \leq 0.4904$$

**(First octant!)**



Because the solar mixing angle does not start as maximal (like in BM mixing) there is a larger region of parameter space in which the reactor and solar mixing angle constraints can be satisfied. This also happens for GR1, GR2, and Hexagonal Mixing. Thus, let this case serve as a representative for them. **Why do I keep mentioning unitarity?**

# The Elephant in the Room



See **1912.10139**  
for details  
describing how to  
generate plot.

Solid lines are when original sum rule is applied with 3 additional sum rules correlating experimental input.

Dashed lines are obtained from treating the experimental distributions as uncorrelated inputs, i.e., not properly using unitarity constraints.

# Conclusion

- The question of why particles have the masses and mixings that they do still remains unsolved, i.e., the flavor problem. However, both current and future experiments are beginning to shine light on possible solutions.
- By assuming a well-known starting point for  $U_\nu$  it is possible to analyze the phenomenological predictions of this starting point by applying the unitary matrix  $U_e$  (see **1801.06377** for the double-rotation cases and more details of the single rotation cases).
- This additional charged lepton rotation gives rise to a nonzero reactor mixing angle and sum rules which allow for correlations between parameters. Perhaps the most important of these correlations are the correlations between the atmospheric and reactor angles.
- Furthermore, even if atmospheric constraints are satisfied in such a simple model, the results for the solar mixing and CP-violating parameter  $\delta$  will further separate these scenarios.
- Studies such as **1801.06377** and **1912.10139** highlight that with the anticipated improvements and measurements of lepton mixing, we may be on the verge of making great progress in understanding the flavor problem.