

# Phenomenology 2021 Symposium



Contribution ID: 1395

Type: BSM

## Explaining $g_\mu - 2$ and $R_{K^{(*)}}$ using the light mediators of $U(1)_{T3R}$

Tuesday, 25 May 2021 15:30 (15 minutes)

Scenarios in which right-handed light Standard Model fermions couple to a new gauge group,  $U(1)_{T3R}$  can naturally generate a sub-GeV dark matter candidate. But such models necessarily have large couplings to the Standard Model, generally yielding tight experimental constraints. We show that the contributions to  $g_\mu - 2$  from the dark photon and dark Higgs largely cancel out in the narrow window where all the experimental constraints are satisfied, leaving a net correction which is consistent with recent measurements from Fermilab. These models inherently violate lepton universality, and UV completions of these models can include quark flavor violation which can explain  $R_{K^{(*)}}$  anomalies as observed at the LHCb experiment after satisfying the  $B_s \rightarrow \mu\mu$  constraint in the allowed parameter space of the model. This scenario can be probed by FASER, SeaQuest, SHIP, LHCb, Belle etc.

### Summary

**Primary author:** GHOSH, Sumit (Texas A&M University)

**Co-authors:** DUTTA, Bhaskar (Texas A&M University); HUANG, Peisi (University fo Nebraska); KUMAR, Jason (University of Hawaii)

**Presenter:** GHOSH, Sumit (Texas A&M University)

**Session Classification:** Flavor III