



Canadian Association  
of Physicists

Association canadienne  
des physiciens et physiciennes

Contribution ID: 212 Type: **Oral Competition (Undergraduate Student) / Compétition orale (Étudiant(e) du 1er cycle)**

## **(U\*) Ab initio calculations of electric dipole moments of light nuclei**

*Thursday 10 June 2021 13:05 (10 minutes)*

In any finite system, the presence of a non-zero permanent electric dipole moment (EDM) would require both parity (P) and time-reversal (T) violation. The standard model predicts a very small CP violation and consequently any observation of the EDM would imply physics beyond the standard model. Thus, EDMs have long been proposed as a way to test these fundamental symmetries. Experimental studies have placed upper bounds on neutron, nuclear and atomic EDMs, while theoretical studies have calculated their magnitudes using a variety of methods. In particular, it has been found that nuclear structure in certain nuclei can enhance the EDM. Here, we use the *ab initio* no-core shell model (NCSM) framework to theoretically investigate the magnitude of the nuclear EDM. We calculate EDMs of several light nuclei using chiral two- and three-body interactions and a PT-violating Hamiltonian based on a one-meson-exchange model. We present a benchmark calculation for  ${}^3\text{He}$ , as well as results for the more complex nuclei  ${}^6,7\text{Li}$ ,  ${}^9\text{Be}$ ,  ${}^{10,11}\text{B}$ ,  ${}^{13}\text{C}$ ,  ${}^{14,15}\text{N}$ , and  ${}^{19}\text{F}$ . Our results suggest that different nuclei can be used to probe different terms of the PT violating interaction. These calculations allow us to suggest which nuclei may be good candidates in the search for a measurable permanent dipole moment.

**Primary authors:** FROESE, Paul (TRIUMF); NAVRATIL, Petr (TRIUMF)

**Presenter:** FROESE, Paul (TRIUMF)

**Session Classification:** R2-5 Theory III (DNP) / Théorie III (DPN)

**Track Classification:** Nuclear Physics / Physique nucléaire (DNP-DPN)