



Canadian Association
of Physicists

Association canadienne
des physiciens et physiciennes

Contribution ID: 217

Type: Oral (Non-Student) / Orale (non-étudiant(e))

Spectrally accurate global-local gyrokinetic simulations of turbulence in tokamak plasmas

Tuesday, 8 June 2021 12:30 (15 minutes)

The suppression of turbulence in fusion plasmas, crucial to the success of next-generation tokamaks such as ITER, depends on a variety of physical mechanisms including the shearing of turbulent eddies via zonal flow and possibly the generation of intrinsic rotation. The turbulence exhibits interesting features such as avalanche structures and self-organisation, and its absence is associated with the formation of internal transport barriers. In order to successfully capture all of these effects in gyrokinetic simulation, one may need to allow for the inclusion of global effects (such as radial profile variation), as well as other often-neglected effects that are small in ρ_* . A careful numerical treatment is necessary to ensure that both the global and local physics are calculated accurately at reasonable expense.

To that end, we develop a novel approach to gyrokinetics where multiple flux-tube simulations are coupled together in a way that consistently incorporates global profile variation while allowing the use of Fourier basis functions, thus retaining spectral accuracy. By doing so, the need for Dirichlet boundary conditions typically employed in global gyrokinetic simulation, where fluctuations are nullified at the simulation boundaries, is obviated. This results in a smooth convergence to the local periodic limit as $\rho_* \rightarrow 0$. In addition, our scale-separated approach allows the use of transport-averaged sources and sinks, offering a more physically motivated alternative to the standard sources based on Krook-type operators.

Having implemented this approach in the flux-tube code `stella`, we study the role of transport barriers and avalanche formation in the transition region between the quiescent core and the turbulent pedestal, as well as the efficacy of intrinsic momentum generation by radial profile variation. Finally, we show that near-marginal plasmas can exhibit a radially localized Dimits shift, where strong coherent zonal flows give way to flows which are more turbulent and smaller scale.

Primary authors: ST-ONGE, Denis (University of Oxford); Prof. PARRA, F. I. (University of Oxford); Prof. BARNES, M. (University of Oxford)

Presenter: ST-ONGE, Denis (University of Oxford)

Session Classification: TS-3 Plasma Physics Symposium (DPP) / Symposium de physique des plasmas (DPP)

Track Classification: Symposia Day (DPP) - Low temperature plasmas/Fusion plasmas (magnetic and inertial confinement)/ Laser plasmas/Basic plasmas