



Canadian Association
of Physicists

Association canadienne
des physiciens et physiciennes

Contribution ID: 364 Type: **Poster Competition (Graduate Student) / Compétition affiches (Étudiant(e) 2e ou 3e cycle)**

(G*) POS-E35 – Novel Methods for Pulse Compression

Wednesday 9 June 2021 13:59 (2 minutes)

Strong field physics, femtosecond ($1 \text{ fs} = 10^{-15}$), and attosecond ($1 \text{ as} = 10^{-18}$) science require intense ultrashort laser pulses for excitation and measurement. Although fs pulses can be generated directly from a laser amplifier, durations necessary for ultrafast science require spectral broadening and compression. The most common spectral broadening technique exploits the nonlinear interaction of intense pulses focused into gas-filled hollow-core fibres [1]. These fibres allow for a long interaction length to generate octave-spanning spectra while maintaining a near Gaussian spatial distribution. However, these fibres can be 1–3 metres on an optical table with additional 1 m focusing and collimating optics, and require continuous backfilled gas flow. More recently, the self-phase modulation in thin crystals broadens the spectrum and uses a self-focusing relay to maintain the beam quality [2]. A combination of solids and gases to generate a supercontinuum has been used to generate attosecond extreme ultraviolet (XUV) pulses [3]. Ultrashort pulses can have a wide variety of applications, including disrupting biological processes for medical procedures, observation of electronic processes on an attosecond scale, and generating XUV and soft X-ray pulses via high harmonic generation or terahertz via difference frequency generation [4]. Here, we compare numerically and experimentally the spectral broadening in solvents and crystals to demonstrate the potential for a compact setup for generating octave spanning pulses for ultrafast experiments.

We find that passing a beam through several cm of methanol produces a supercontinuum spectrum that can support few-fs pulses. Similar to the multi-plate thin crystal method, we place a series of 1 cm cuvettes of methanol around two foci in order to reduce self-focussing effects and to avoid filamentation within the material. We compress these pulses using dispersion compensating mirrors, and characterize the resulting pulses using frequency resolved optical gating (FROG) and a novel measurement scheme. Our simulations and models predict a sub-10 fs bandwidth-limited pulse resulting from two subsequent compressions using a total of 4 cm of methanol to generate a supercontinuum. Using this method, experimentally, we have achieved a broad spectrum capable of supporting sub-30 fs pulses.

[1] E. Haddad, R. Safaei, A. Leblanc, R. Piccoli, Y.G. Jeong, H. Ibrahim, B.E. Schmidt, R. Morandotti, L. Razzari, F. Légaré, P. Lassonde. "Molecular gases for pulse compression in hollow core fibers," *Optics Express* (2018).

[2] Y.-C. Cheng, C.-H. Lu, Y.-Y. Lin, and A. H. Kung, "Supercontinuum generation in a multi-plate medium," *Optics Express* (2016).

[3] T. J. Hammond, S. Monchocé, C. Zhang, G. Vampa, D. Klug, A. Y. Naumov, D. M. Villeneuve, and P. B. Corkum "Integrating solids and gases for attosecond pulse generation," *Nature Photonics* (2017).

[4] M. Cheng, A. Reynolds, H. Widgren, and M. Khalil, "Generation of tunable octave-spanning mid-infrared pulses by filamentation in gas media," *Optics Letters* (2012).

Primary author: STEPHEN, Jacob (University of Windsor)

Co-authors: JAYALATH ARACHCHIGE, Chathurangani (University of Windsor); HAMMOND, TJ (University of Windsor)

Presenter: STEPHEN, Jacob (University of Windsor)

Session Classification: W-POS-E #28-40 Poster Session (DAMOFC) / Session d'affiches (DPAMOC)

Track Classification: Atomic, Molecular and Optical Physics, Canada / Physique atomique, moléculaire et photonique, Canada (DAMOFC-DPAMOC)