



Canadian Association  
of Physicists

Association canadienne  
des physiciens et physiciennes

Contribution ID: 301      Type: **Oral not-in-competition (Graduate Student) / Orale non-compétitive (Étudiant(e) du 2e ou 3e cycle)**

## Emergence of light-induced states in the few-photon ionization of atomic helium\*

Monday, 7 June 2021 16:59 (4 minutes)

In this joint experimental and theoretical work [1], photoelectron emission from excited states of laser-dressed atomic helium is analyzed. We successfully demonstrate a method that is complementary to transient absorption (e.g. [2]) for the assignment of light-induced states (LIS). The experiment is carried out at DESY in Hamburg and uses the FLASH2 free-electron laser to produce an extreme ultraviolet (XUV) pulse to which the helium atom is subjected along with a temporally overlapping infrared (IR) pulse in the multi-photon ionization regime ( $\approx 10^{12}$  W/cm<sup>2</sup>). Analysis of the experiment occurs at the reaction microscope (REMI) end station [3] at FLASH2. The XUV pulse is scanned over the energy range 20.4 eV to 24.6 eV, corresponding to excited states of helium. The resonant, electric dipole-allowed  $nP$  states corresponding to a first step of single XUV photon excitation are shown to lead to ionization, independent of whether or not the lasers temporally overlap. However, dipole-forbidden transitions to  $nS$  and  $nD$  states corresponding to multiphoton (XUV  $\pm$   $n$ IR) excitation are observed during temporal overlap. Studying photoelectron angular distributions (PADs) in the case where the ionization pathway of a LIS is difficult to resolve energetically allows for an unambiguous determination of the dominant LIS. The IR intensity and relative polarization between the lasers are varied to control the ionization pathway. Numerical solutions of the time-dependent Schrödinger equation within a single-active electron model with a local potential completely support the experimental findings in this project.

[1] S. Meister *et al.*, Phys. Rev. A **102** (2020) 062809; Phys. Rev. A **103** (2021) in press.

[2] S. Chen *et al.*, Phys. Rev. A **86** (2012) 063408.

[3] S. Meister *et al.*, Applied Sciences **10** (2020) 2953.

\*work supported by NSERC, NSF, and XSEDE

**Primary authors:** BONDY, Aaron; MEISTER, Severin (Max-Planck-Institute for Nuclear Physics); SCHNORR, Kirsten (Paul Scherrer Institute); AUGUSTIN, Sven (Paul Scherrer Institute); LINDENBLATT, Hannes (Max-Planck-Institute for Nuclear Physics); TROST, Florian (Max-Planck-Institute); XIE, Xinhua (Paul Scherrer Institute); BRAUNE, Markus (Deutsches Elektronen-Synchrotron); TREUSCH, Rolf (Deutsches Elektronen-Synchrotron); MANSCHWETUS, Bastian (Deutsches Elektronen-Synchrotron); SCHIRMEL, Nora (Deutsches Elektronen-Synchrotron); REDLIN, Harald (Deutsches Elektronen-Synchrotron); DOUGUET, Nicolas (Kennesaw State University); PFEIFER, Thomas (Max-Planck-Institute for Nuclear Physics); BARTSCHAT, Klaus (Drake University); MOSHAMMER, Robert (Max-Planck-Institute for Nuclear Physics)

**Presenter:** BONDY, Aaron

**Session Classification:** M4-1 Optical spectroscopy (DAMOPC) / Spectroscopie optique (DPAMPC)

**Track Classification:** Atomic, Molecular and Optical Physics, Canada / Physique atomique, moléculaire et photonique, Canada (DAMOPC-DPAMPC)