



Contribution ID: 205

Type: Theory talk

## Thermalization time constrained by high-pt QGP tomography

Friday, 21 May 2021 10:10 (20 minutes)

We show that high- $p_{\perp}$   $R_{AA}$  and  $v_2$  are way more sensitive to the QGP thermalization time,  $\tau_0$ , than the distributions of low- $p_{\perp}$  particles, and that the high- $p_{\perp}$  observables prefer relatively late thermalization at  $\tau_0 \sim 1$  fm/c. To calculate high- $p_{\perp}$   $R_{AA}$  and  $v_2$ , we employ our newly developed DREENA-A formalism, which combines state-of-the-art dynamical energy loss model with 3+1 dimensional hydrodynamical simulations. The model applies to both light and heavy flavor, and we predict a larger sensitivity of heavy observables to the thermalization time. Elliptic flow parameter  $v_2$  is also more sensitive to  $\tau_0$  than  $R_{AA}$  due to non-trivial differences in the evolution of in-plane and out-of-plane temperature profiles. This presents the first time when a parameter describing bulk QGP has been constrained by high- $p_{\perp}$  observables and related theory, i.e., by so-called QGP tomography.

### Collaboration

**Authors:** Mr STOJKU, Stefan (Institute of Physics University of Belgrade); Dr AUVINEN, Jussi (Institute of Physics University of Belgrade); Prof. DJORDJEVIC, Marko (University of Belgrade Faculty of Biology); Dr HUOVINEN, Pasi (Institute of Physics University of Belgrade); Dr DJORDJEVIC, Magdalena (Institute of Physics University of Belgrade)

**Presenter:** Mr STOJKU, Stefan (Institute of Physics University of Belgrade)

**Session Classification:** Bulk (Small systems)

**Track Classification:** Other