



✂ Particle density evolution in the expanding universe

At classical level, number density n_{i_k} of any particle species included in $\{i\} \leftrightarrow \{f\}$ reactions,

$$\dot{n}_{i_k} + 3\mathcal{H}n_{i_k} = -\hat{\gamma}_{fi} + \hat{\gamma}_{if} + \dots \quad (1)$$

Reaction rates

$$\hat{\gamma}_{fi} = \int \prod_{\{i\}} [d\mathbf{p}_i] \hat{f}_i(p_i) \int \prod_{\{f\}} [d\mathbf{p}_f] (2\pi)^4 \delta^{(4)}(p_f - p_i) |\dot{M}_{fi}|^2 \equiv \int_{i \rightarrow f} \hat{f}_i |\dot{M}_{fi}|^2 \quad (2)$$

← classical phase-space densities \hat{f}_i (Maxwell-Boltzmann densities in equilibrium),

← zero-temperature Feynman rules entering the calculation of \dot{M}_{fi} .

✂ CP asymmetries at zero-temperature

The S -matrix unitarity $S^\dagger = S^{-1}$, or $1 - iT^\dagger = (1 + iT)^{-1}$, for the squared amplitude leads to

$$iT^\dagger = iT - (iT)^2 + (iT)^3 - \dots \Rightarrow |T_{fi}|^2 = -iT_{if}iT_{fi} + \sum_n iT_{in}iT_{nf}iT_{fi} - \dots \quad (3)$$

Divide by $V_4 = (2\pi)^4 \delta^{(4)}(0)$ to obtain $(2\pi)^4 \delta^{(4)}(p_f - p_i) |\dot{M}_{fi}|^2$. In CPT symmetric theory, the CP asymmetry $\Delta|T_{fi}|^2 = |T_{fi}|^2 - |T_{if}|^2$ can be expressed as [1]

$$\begin{aligned} \Delta|T_{fi}|^2 &= \sum_n (iT_{in}iT_{nf}iT_{fi} - iT_{if}iT_{fn}iT_{ni}) \\ &\quad - \sum_{n,m} (iT_{in}iT_{nm}iT_{mf}iT_{fi} - iT_{if}iT_{fm}iT_{mn}iT_{ni}) \\ &\quad + \dots \end{aligned} \quad (4)$$

From Eq. (4) the CPT and unitarity relations [2–4]

$$\sum_f \Delta|T_{fi}|^2 = 0 \quad (5)$$

are manifest at any perturbative order [1].

✂ Thermal effects at the lowest order

As an example we consider $N_i \rightarrow lH$ decay within the seesaw type-I model. Right-handed neutrinos N_i , standard model lepton l and Higgs doublets H interact via Yukawa interactions

$$\mathcal{L} \supset -\frac{1}{2} \bar{M}_i \bar{N}_i N_i - \left(\mathcal{Y}_{\alpha i} \bar{N}_i P_L l_\alpha H + \text{H.c.} \right). \quad (6)$$

To the lowest $\mathcal{O}(\mathcal{Y}^2)$ order,

$$\hat{\gamma}_{N_i \rightarrow lH} = \int_{N_i \rightarrow lH} \hat{f}_{N_i} |\dot{M}|^2, \quad |\dot{M}|^2 = 4p_{N_i} p_l \sum_\alpha |\mathcal{Y}_{\alpha i}|^2 \quad (7)$$

that can be represented by the cut of the forward diagram Fig. 1a. Diagram in Fig. 1b leads to spurious contribution, equivalent to $\hat{\gamma}_{N_i \rightarrow lH}$ times the total number of Higgs particles in the universe. Clearly, such contributions have to be omitted. Diagram in Fig. 1c corresponds to

$$\hat{\gamma}_{N_i H \rightarrow lHH} = \int_{N_i \rightarrow lH} \hat{f}_{N_i} \hat{f}_H |\dot{M}|^2 \quad (8)$$

with the same $|\dot{M}|^2$ as in Eq. (7). To sum up the contributions such as in Fig. 1c avoiding those analogous to Fig. 1b, we shall draw the forward diagrams on a cylindrical surface [5]. In thermal equilibrium $\hat{f}_H = e^{-E/T}$ and the summation leads to

$$\sum_{w=0}^{\infty} \hat{f}_H^w = \frac{e^{E/T}}{e^{E/T} - 1} \equiv 1 + f_H^{\text{eq}}. \quad (9)$$




Figure 1: Planar and cylindrical diagrams contributing to the $N_i \rightarrow lH$ zero-temperature reaction rate and its thermal corrections.

Similarly, with alternating signs due to the crossing of fermionic legs, the Fermi-Dirac distributions for N_i and l are obtained. Thus, we introduce

$$\gamma_{N_i \rightarrow lH} = \int_{N_i \rightarrow lH} f_{N_i} (1 + f_H^{\text{eq}}) (1 - f_l^{\text{eq}}) |\dot{M}|^2. \quad (10)$$

✂ Thermal corrections to the lepton number source term




Figure 2: Lepton number violating contributions to the CP asymmetry in $N_i \rightarrow lH$ decay (Figs. 2a and 2b) and the s -channel part of the real-intermediate-state-subtracted $lH \rightarrow \bar{l}\bar{H}$ scattering (Fig. 2c). The latter can be obtained by a cyclic permutation of the diagrams in Fig. 2a using the approach of Refs. [1, 6].

In leptogenesis, the matter asymmetry is generated from the $N_i \rightarrow lH$ decays and $lH \rightarrow \bar{l}\bar{H}$ scatterings (real-intermediate-state-subtracted scatterings). The zero-temperature asymmetry of the squared amplitude for the N_i decay, before the momentum integration is performed, equals

$$\Delta|\dot{M}|^2 = 4p_l p_{\bar{l}} \sum_{j \neq i} 2\text{Im} \left[\left(\mathcal{Y} \mathcal{Y}^\dagger \right)_{ij}^2 \right] \left(\frac{2M_i M_j}{M_i^2 - M_j^2} + \frac{M_i M_j}{(p_l - p_{\bar{H}})^2 - M_j^2} \right). \quad (11)$$

At finite temperature, the on-shell parts of the propagators are modified, affecting the reaction rates asymmetries. If, for example, the first term in

$$\frac{\bar{H}}{N_j} + \frac{\bar{H}}{N_j} + \frac{\bar{H}}{N_j} + \dots, \quad (12)$$

occurs as a subdiagram, depending on the overall kinematics, the positive frequency on-shell part of the \bar{H} propagator may contribute or not. If it does, all the other terms do as well. Thus, the \bar{H} line can be replaced by

$$\frac{i}{p_H^2 + i\epsilon} + 2\pi \sum_{w=1}^{\infty} \hat{f}_H^w \theta(p_H^0) \delta(p_H^2) = \frac{i}{p_H^2 + i\epsilon} + 2\pi f_H^{\text{eq}} \theta(p_H^0) \delta(p_H^2) \quad (13)$$

corresponding to the usual form of the positive frequency part of a thermal propagator. The negative frequencies contribute through the diagrams with opened \bar{H} legs flipped to the opposite side.

Therefore, to take the thermal corrections to the reaction rate asymmetries into account, we should consider all the windings of the propagators involved, leading to

$$\Delta\gamma_{N_i \rightarrow lH} = \int_{N_i \rightarrow lH \rightarrow \bar{l}\bar{H}} f_{N_i} (1 - f_l^{\text{eq}}) (1 + f_H^{\text{eq}}) (1 - f_{\bar{l}}^{\text{eq}}) (1 + f_{\bar{H}}^{\text{eq}}) \Delta|\dot{M}|^2, \quad (14a)$$

$$\Delta\gamma_{\bar{l}\bar{H} \rightarrow N_i} = \int_{N_i \rightarrow lH \rightarrow \bar{l}\bar{H}} f_{\bar{l}}^{\text{eq}} f_{\bar{H}}^{\text{eq}} (1 - f_{N_i}) (1 - f_l^{\text{eq}}) (1 + f_H^{\text{eq}}) \Delta|\dot{M}|^2, \quad (14b)$$

$$\Delta\gamma_{lH \rightarrow \bar{l}\bar{H}} = \int_{N_i \rightarrow lH \rightarrow \bar{l}\bar{H}} f_l^{\text{eq}} f_H^{\text{eq}} (1 - f_{\bar{l}}^{\text{eq}}) (1 + f_{\bar{H}}^{\text{eq}}) (1 - f_{N_i}) \Delta|\dot{M}|^2. \quad (14c)$$

Using the detailed balance conditions, such as

$$f_{N_i}^{\text{eq}} (1 - f_l^{\text{eq}}) (1 + f_H^{\text{eq}}) = (1 - f_{N_i}^{\text{eq}}) f_l^{\text{eq}} f_H^{\text{eq}}, \quad (15)$$

and unitarity relations from Eq. (5) for asymmetries in Eq. (14), we obtain for the lepton number evolution

$$\dot{n}_L + 3\mathcal{H}n_L = \int_{N_i \rightarrow lH \rightarrow \bar{l}\bar{H}} \delta f_{N_i} (1 - f_l^{\text{eq}}) (1 + f_H^{\text{eq}}) (1 - f_{\bar{l}}^{\text{eq}} + f_{\bar{H}}^{\text{eq}}) \Delta|\dot{M}|^2 + \text{wash-out terms}, \quad (16)$$

where $n_L \equiv n_l - n_{\bar{l}}$. Remarkably, the result is in agreement with non-equilibrium QFT calculations [7–9]. The source term vanishes in equilibrium, when $\delta f_{N_i} = f_{N_i} - f_{N_i}^{\text{eq}} = 0$.

✂ Conclusions

- ← **A diagrammatic concept connecting the classical Boltzmann equation and quantum kinetic theory has been introduced.**
- ← **Statistical factors due to on-shell intermediate states have been formally represented by the cuts of forward diagrams with multiple spectator lines.**
- ← **Results are in agreement with the direct closed-time-path derivation of non-equilibrium quantum field theory [7–9].**
- ← **Unitarity and CPT relations between the reaction rate asymmetries can be derived in our approach to kinetic theory in the same way as in $T = 0$ calculations.**

References

- [1] T. Blažek and P. Maták, *CP Asymmetries and Higher-Order Unitarity Relations*, *Phys. Rev. D* **103** (2021) [2102.05914].
- [2] E.W. Kolb and S. Wolfram, *Baryon Number Generation in the Early Universe*, *Nucl. Phys. B* **172** (1980) 224.
- [3] A. Hook, *Unitarity constraints on asymmetric freeze-in*, *Phys. Rev. D* **84** (2011) 055003.
- [4] I. Baldes, N.F. Bell, K. Petraki and R.R. Volkas, *Particle-antiparticle asymmetries from annihilations*, *Phys. Rev. Lett.* **113** (2014) 181601.
- [5] T. Blažek and P. Maták, *Cutting Rules on Cylinder: Bottom-up Approach to Quantum Kinetic Theory*, **2104.06395**.
- [6] E. Roulet, L. Covi and F. Vissani, *On the CP asymmetries in Majorana neutrino decays*, *Phys. Lett. B* **424** (1998) 101 [hep-ph/9712468].
- [7] M. Garny, A. Hohenegger, A. Kartavtsev and M. Lindner, *Systematic approach to leptogenesis in nonequilibrium QFT: Vertex contribution to the CP-violating parameter*, *Phys. Rev. D* **80** (2009) 125027 [0909.1559].
- [8] M. Garny, A. Hohenegger, A. Kartavtsev and M. Lindner, *Systematic approach to leptogenesis in nonequilibrium QFT: Self-energy contribution to the CP-violating parameter*, *Phys. Rev. D* **81** (2010) 085027 [0911.4122].
- [9] B. Garbrecht, *Leptogenesis: The Other Cuts*, *Nucl. Phys. B* **847** (2011) 350 [1011.3122].