

Reconstruction of b - and c - jets at e^+e^- Higgs Factories with ParticleFlow detectors

International Workshop on Future Linear Colliders, LCWS2021

Yasser Radkhorrami^{1,2}, Jenny List¹

¹DESY, Hamburg

²Universität Hamburg, Hamburg

March 18, 2021



CLUSTER OF EXCELLENCE
QUANTUM UNIVERSE


HELMHOLTZ
RESEARCH FOR GRAND CHALLENGES




Higgs production mechanisms and decay modes at e^+e^- colliders

- Higgs strahlung is dominant Higgs production mechanism at 250 GeV


$P(e^-, e^+) = (-0.8, 0.3)$, $M_h = 125$ GeV



- Most frequent Higgs decay mode: $H \rightarrow b\bar{b}$




- Extremely challenging in Hadron colliders: $H \rightarrow c\bar{c}$





Semi-leptonic b / c decays

- Number of B-/C-hadron semileptonic decays (SLD) in $e^+e^- \rightarrow b\bar{b}$ events

nBSLD				
	0	1	2	
nCSLD	0	34%	24%	4%
1	18%	12%	2%	
2	3%	2%	0%	



- Mis-reconstruction of $b\bar{b}$ invariant mass due to missing neutrino energy from semi-leptonic decays
- Can the missing momentum be retrieved from event and decay kinematics?




Concept of ν -correction in a semi-leptonic decay


- ▶ Find heavy-quark jets: Identify b or c jet \rightarrow flavour tag
- ▶ Find semi-leptonic decay(s): Identify lepton in jet if present
 \rightarrow possible using detector's high granularity
- ▶ Estimate neutrino energy from decay kinematics:
 - ▶ Assign B^0 or D^0 meson mass to mother hadron.
 - ▶ Reconstruct flight direction of mother hadron from position of primary and secondary vertex.
 - ▶ Calculate neutrino momentum: up to 2-fold ambiguity.
- ▶ As proof-of-principle: CHEAT from MC truth

The neutrino momentum can be determined up to a two-fold ambiguity

Can we use overall event kinematics to decide between solutions? \Rightarrow kinematic fit!




- ▶ Closure test: fully cheated information ($e^+e^- \rightarrow b\bar{b}$ at $\sqrt{s} = 500$ GeV)



Kinematic fit

- Kinematic fit: adjustment of measured quantities under certain kinematic constraints:
 - ▶ Energy and momentum conservation
 - ▶ Invariant masses of particles



Exploit well-known initial state in e^+e^- colliders

⇒ need error parametrization, in particular for jets

- Minimize χ^2 :

$$\chi^2(a, \xi, f) = (\eta - a)^T V^{-1}(\eta - a) - 2\lambda^T f(a, \xi)$$

η : vector of measured kinematic variables (x)

a : vector of fitted quantities

ξ : vector of unmeasured kinematic variables

V : **covariance matrix**

λ : Lagrange multipliers

$f(a, \xi)$: vector of constraints

Jet specific energy resolution

Parametrize sources of uncertainties (assumed uncorrelated) in jet energy measurements (ErrorFlow):

$$\sigma_{E_{jet}} = \sigma_{Det} \oplus \sigma_{Conf} \oplus \sigma_{\nu} \oplus \sigma_{Clus} \oplus \sigma_{Had}$$

- ▶ σ_{Det} : Detector resolution using track and cluster parameters
- ▶ σ_{Conf} : Particle confusion in Particle Flow Algorithm
Estimated based on jet energy and neutral hadron / photon energy fractions
- ▶ σ_{ν} : Semi-leptonic decays: error propagation from neutrino correction
currently none since cheating
- ▶ σ_{Clus} : Misassignment of particles in the jet clustering, has not been included yet
- ▶ σ_{Had} : Mismodeling of QCD effects in parton shower and hadronization, has not been included yet

DESY-THESIS-2017-045




ErrorFlow: Jet Error Parametrisation from Particle Flow Objects (PFO)

Energy

Error estimation in PFO level:



- ▶ **Photons**: energy error is perfectly modeled.
($\sigma \sim 1$)
- ▶ **Charged PFOs**: uncertainties propagated from track fit covariance matrix
 - ▶ uncertainties 30% too small
 - ▶ possible future improvement from track refitting with specific mass hypothesis after particle ID
- ▶ **Neutral Hadrons**: energy and energy error are significantly overestimated.
work on improvement in progress.



ErrorFlow: Jet Error Parametrisation from Particle Flow Objects (PFO)

Angles

The angular uncertainties obtained directly from track parameters / cluster position errors




⇒ Scale σ_θ and σ_ϕ by factor ~ 1.3 (for photons) and ~ 1.8 (for neutral hadrons)



Uncertainties in jet-level: Energy

Propagation of errors from PFOs to jets:



- ▶ Transform the covariance matrix of each PFO (E, x, y, z for clusters, track parameters for charged) to (E, p_x, p_y, p_z)
- ▶ Add up covariance matrices of all PFOs
- ▶ Add confusion term for jet energy
 - ▶ calculate using jet energy composition
- ▶ Transform to (E, θ, ϕ, m)



Confusion term improves the estimate of the jet energy uncertainty, but not quite enough ⇒ need adjustment
 ⇒ use scaling factor 1.2 in Kinematic fit



Uncertainties in jet-level: θ & ϕ



Jet angular uncertainties need scaling factor ~ 1.6



Application of kinematic fit to $e^+e^- \rightarrow ZH \rightarrow \mu\bar{\mu} b\bar{b}$ events

Parameters of jets and leptons are varied within their uncertainties to satisfy 5 constraints:
 Conservation of momentum (hard constraints):

- ▶ p_x : e^+e^- crossing angle: 14 mrad

$$\Sigma p_x = \sqrt{s} \times \sin 0.007 \approx 1.75 \text{ GeV}$$

- ▶ p_y : $\Sigma p_y = 0$

- ▶ p_z : $\Sigma p_z = 0$


arXiv:0901.4656

Conservation of total energy (hard constraint):

$$\nabla E_{lab} = 2\sqrt{\left(\frac{\sqrt{s}}{2}\right)^2 + (\Sigma p_x)^2}$$

Constrain di-muon mass to agree with m_Z within its natural width
 (soft constraint):


$$\nabla m_Z = 91.2 \text{ GeV}, \sigma_{m_Z} = \frac{2.4952}{2}$$




Kinematic fit performance in $e^+e^- \rightarrow ZH \rightarrow \mu\bar{\mu} b\bar{b}$ at $\sqrt{s} = 250$ GeV

without semi-leptonic decays

► pull distribution



► fit probability





Improved kinematic fit performance with full CovMat of jets + scaled jet energy uncertainty



Kinematic fit performance in $e^+e^- \rightarrow ZH \rightarrow \mu\bar{\mu}bb$ at $\sqrt{s} = 250$ GeV (cntd.)

without semi-leptonic decays





Improved kinematic fit performance with full CovMat of jets + scaled jet angular uncertainties



Higgs mass in presence of SLDs

ν -correction and kinematic fit on $H \rightarrow b\bar{b}$



- ▶ Applying **kinematic fit and ν -correction** gives huge improvement on Higgs mass reconstruction



Conclusions

- ▶ Higgs mass reconstruction essential eg in ZZH vs ZHH separation (Higgs self-coupling measurement)
- ▶ Heavy flavour jets are essential for Higgs physics
- ▶ Correction of semi-leptonic decays of heavy flavour jets is important for Higgs mass reconstruction
 - ▶ Neutrino momentum can be reconstructed up to a sign ambiguity
 - ▶ Ambiguity can be resolved by kinematic fit
 - ▶ Next: remove the partial cheating from the neutrino correction
- ▶ Kinematic fit exploits well-known initial state in e^+e^- colliders and requires excellent understanding of jet measurement
- ▶ ILD as a Particle Flow detector provides full detail for estimating jet measurement uncertainties


BACKUP




International Large Detector (ILD)

arXiv:2003.01116

Momentum Resolution




Jet Energy Resolution ($E_{PFO} + E_\nu^{MC}$)




International Large Detector (ILD)

arXiv:2003.01116

Impact Parameter Resolution, d_0



Impact Parameter Resolution, z_0




Concept of ν -correction in a semi-leptonic decay


- ▶ Find heavy-quark jets: Identify b or c jet \rightarrow flavour tag
- ▶ Find semi-leptonic decay(s): Identify lepton in jet if present \rightarrow possible using detector's high granularity
- ▶ Estimate neutrino energy from decay kinematics:
 - ▶ Assign B^0 or D^0 meson mass to mother hadron.
 - ▶ Reconstruct flight direction of mother hadron from position of primary and secondary vertex.
 - ▶ Calculate neutrino momentum: up to 3-fold ambiguity.
- ▶ As proof-of-principle: CHEAT from MC truth
 - ▶ Lepton ID
 - ▶ Flavour tag
 - ▶ Mother hadron mass
 - ▶ **Associate of reconstructed particles to secondary vertex**
 - ▶ Momenta of visible decay products

The neutrino momentum can be determined up to a two-fold ambiguity

Can we use overall event kinematics to decide between solutions? \Rightarrow kinematic fit!




- ▶ Closure test: fully cheated information ($e^+e^- \rightarrow b\bar{b}$ at $\sqrt{s} = 500$ GeV)



correcting neutrino energy

4-vector based approach

- (E, \vec{p})-based approach



$$\vec{p}_{\nu, \perp} = -\vec{p}_{vis, \perp}$$

$$\vec{p}\nu, \parallel = \frac{1}{2D}(-A \pm \sqrt{A^2 - BD})\hat{n}$$

$$A = p_{vis, \parallel}(2p_{vis, \perp}^2 + m_{vis}^2 - m_X^2)$$


$$B = 4p_{vis, \perp}^2 E_{vis}^2 - (2p_{vis, \perp}^2 + m_{vis}^2 - m_X^2)^2$$

$$D = E_{vis}^2 - p_{vis, \parallel}^2$$

$$\hat{n} = \frac{\vec{p}_{vis, \parallel}}{|\vec{p}_{vis, \parallel}|}$$

The neutrino momentum can be determined up to a two-fold ambiguity

- closure test: apply correction with fully cheated information and compare with true neutrino energy




Correcting neutrino energy

Rapidity based approach

Rapidity under Lorentz-transformations \sim velocity under Galileo-transformations: $\omega = \omega_X + \omega'$; $\omega = \frac{1}{2} \ln \frac{E+p'_\parallel}{E-p'_\parallel}$

ω : rapidity in lab frame , ω' : rapidity in rest frame of X , ω_X : rapidity of X in lab frame



$$E_\nu = E_X - E_{vis}$$

$$E_X = \frac{E_{vis}' E_{vis}' - p_{vis\parallel} p_{vis\perp}'}{m_{vis}^2 + p_{vis\perp}^2} m_X$$


$$E'_{vis} = \frac{m_X^2 + m_{vis}^2}{2m_X}$$

$$p'_{vis\parallel} = \pm \sqrt{\left(\frac{m_X^2 - m_{vis}^2}{2m_X}\right)^2 - p_{vis\perp}^2}$$

The neutrino momentum can be determined up to a two-fold ambiguity

Can we use overall event kinematics to decide between solutions? \Rightarrow kinematic fit!

- ▶ Closure test: fully cheated information
 $(e^+e^- \rightarrow b\bar{b}$ at $\sqrt{s} = 500$ GeV)



Event selection

Select $e^+e^- \rightarrow ZH \rightarrow \mu\bar{\mu}b\bar{b}$ events at $\sqrt{s} = 250$ GeV with (exactly) 2-leptons + 2-jets final state:


► IsolatedLeptonTagging

Training for the IDR 500 GeV samples is used,

1. Lepton ID: μ^\pm
Deposited energy in subdetectors
2. Vertex: primary or secondary
Significance of impact parameters (d_0 , z_0)
3. Isolated: not belong to jets

► FastJetProcessor

- Exclusive k_t (Durham) algorithm (no overlay)
 - Find smallest of (d_{ij}, d_{iB})
- $$d_{ij} = 2 \min(E_i^2, E_j^2)(1 - \cos \theta_{ij})$$
- i, j : particles, B : Beam
- $d_{ij} < d_{iB}$: combine $i \& j$ as pseudojet(p): $p_i + p_j$
 - $d_{iB} < d_{ij}$: remove particle i from list
 - Repeat iteration until d_{ij} or $d_{iB} > d_{cut}$ (threshold)




IsolatedLeptonTagging has not been trained for new software at 250 GeV yet!



event selection


separate Higgs decay modes: $H \rightarrow b\bar{b}$, cheat from MCTruth



$\frac{2}{3}$ of $b\bar{b}$ jets contain at-least one semi-leptonic decay \Rightarrow Frequent $H \rightarrow b\bar{b}$ needs neutrino correction.



Neutral PFO identification by Pandora



Majority of identified photons are true photons.

No explicit decision for mass of identified neutral hadrons due to their multiplicity.



Pandora treatment with Neutral Hadrons

What Pandora does:

- ▶ Cluster energy is assigned to PFO(massless) energy
 $E_{PFO} = |\vec{p}_{PFO}| = E_{cluster}$
- ▶ Neutral Hadrons are identified as neutron
- ▶ neutron mass is set for PFO \Rightarrow **inconsistent 4-momentum!**
- ▶ CovMat of Neutral PFO is calculated (using inconsistent 4-momentum):
 $\text{CovMat}(\vec{p}, E) = J^T \text{CovMat}(\vec{x}_{clu}, E_{clu}) J$

$$J = \begin{pmatrix} \frac{\partial p_x}{\partial x_c} & \frac{\partial p_y}{\partial x_c} & \frac{\partial p_z}{\partial x_c} & \frac{\partial E}{\partial x_c} \\ \frac{\partial p_x}{\partial p_y} & \frac{\partial p_y}{\partial p_y} & \frac{\partial p_z}{\partial p_y} & \frac{\partial E}{\partial p_y} \\ \frac{\partial p_x}{\partial y_c} & \frac{\partial p_y}{\partial y_c} & \frac{\partial p_z}{\partial y_c} & \frac{\partial E}{\partial y_c} \\ \frac{\partial p_x}{\partial z_c} & \frac{\partial p_y}{\partial z_c} & \frac{\partial p_z}{\partial z_c} & \frac{\partial E}{\partial z_c} \\ \frac{\partial p_x}{\partial E_c} & \frac{\partial p_y}{\partial E_c} & \frac{\partial p_z}{\partial E_c} & \frac{\partial E}{\partial E_c} \end{pmatrix}$$

$\text{CovMat}(\vec{p}, E)$ of Neutral PFOs depend on the mass assumption.

Suggestion: Take consistent 4-momentum of massive neutral hadrons for CovMat calculations.



CovMat of Neutral PFOs

- ▶ Current CovMat calculation (MarlinReco/Analysis/AddClusterProperties)

$$E_{PFO} = |\vec{p}_{PFO}| = E_{clu}, p_x = E_{clu} \frac{x}{r}, p_y = E_{clu} \frac{y}{r}, p_z = E_{clu} \frac{z}{r}$$

- ▶ Alternative CovMat calculation (taking consistent 4-momentum of neutral hadrons)

$$E_{PFO} = \sqrt{|\vec{p}_{PFO}|^2 + m_{PFO}^2} = \sqrt{E_{clu}^2 + m_n^2}$$

$$J = \begin{pmatrix} E_{clu} \frac{r^2-x^2}{r^3} & -E_{clu} \frac{xy}{r^3} & -E_{clu} \frac{xz}{r^3} & 0 \\ -E_{clu} \frac{xy}{r^3} & E_{clu} \frac{r^2-y^2}{r^3} & -E_{clu} \frac{yz}{r^3} & 0 \\ -E_{clu} \frac{xz}{r^3} & -E_{clu} \frac{yz}{r^3} & E_{clu} \frac{r^2-z^2}{r^3} & 0 \\ \frac{x}{r} & \frac{y}{r} & \frac{z}{r} & 1 \end{pmatrix} \rightarrow J = \begin{pmatrix} E_{clu} \frac{r^2-x^2}{r^3} & -E_{clu} \frac{xy}{r^3} & -E_{clu} \frac{xz}{r^3} & 0 \\ -E_{clu} \frac{xy}{r^3} & E_{clu} \frac{r^2-y^2}{r^3} & -E_{clu} \frac{yz}{r^3} & 0 \\ -E_{clu} \frac{xz}{r^3} & -E_{clu} \frac{yz}{r^3} & E_{clu} \frac{r^2-z^2}{r^3} & 0 \\ \frac{E}{E_{clu}} \cdot \frac{x}{r} & \frac{E}{E_{clu}} \cdot \frac{y}{r} & \frac{E}{E_{clu}} \cdot \frac{z}{r} & 1 \end{pmatrix}$$

using error propagation, PFO angular uncertainties are calculated directly from cluster position error:

$$\sigma_\theta^2 = (\frac{\partial\theta}{\partial x})^2 \sigma_x^2 + (\frac{\partial\theta}{\partial y})^2 \sigma_y^2 + (\frac{\partial\theta}{\partial z})^2 \sigma_z^2 + \frac{\partial\theta}{\partial x} \frac{\partial\theta}{\partial y} \sigma_{xy} + \frac{\partial\theta}{\partial x} \frac{\partial\theta}{\partial z} \sigma_{xz} + \frac{\partial\theta}{\partial y} \frac{\partial\theta}{\partial z} \sigma_{yz}$$

$$\sigma_\phi^2 = (\frac{\partial\phi}{\partial x})^2 \sigma_x^2 + (\frac{\partial\phi}{\partial y})^2 \sigma_y^2 + \frac{\partial\phi}{\partial x} \frac{\partial\phi}{\partial y} \sigma_{xy}$$

MUST: angular and energy uncertainties remain unchanged!



CovMat of Jets

- AddClusterProperties/FourMomentumCovMat: $\text{CovMat}(\text{cluster}/\text{track}) \rightarrow \text{CovMat}(\vec{p}, E)$

- Current CovMat calculation (inconsistent 4-momentum of neutral hadrons):

$$E_{PFO} = |\vec{p}_{PFO}| = E_{clu}, p_x = E_{clu} \frac{x}{r}, p_y = E_{clu} \frac{y}{r}, p_z = E_{clu} \frac{z}{r}, m_{PFO} = m_n$$

- Alternative CovMat calculation (taking consistent 4-momentum of neutral hadrons)

$$E_{PFO} = \sqrt{|\vec{p}_{PFO}|^2 + m_{PFO}^2} = \sqrt{E_{clu}^2 + m_n^2}$$

$J_{(wrong)} \rightarrow J_{(right)}$

$$\begin{pmatrix} E_{clu} \frac{r^2 - x^2}{r^3} & -E_{clu} \frac{xy}{r^3} & -E_{clu} \frac{xz}{r^3} & 0 \\ -E_{clu} \frac{xy}{r^3} & E_{clu} \frac{r^2 - y^2}{r^3} & -E_{clu} \frac{yz}{r^3} & 0 \\ -E_{clu} \frac{xz}{r^3} & -E_{clu} \frac{yz}{r^3} & E_{clu} \frac{r^2 - z^2}{r^3} & 0 \\ \frac{x}{r} & \frac{y}{r} & \frac{z}{r} & 1 \end{pmatrix}_{wrong} \rightarrow \begin{pmatrix} E_{clu} \frac{r^2 - x^2}{r^3} & -E_{clu} \frac{xy}{r^3} & -E_{clu} \frac{xz}{r^3} & 0 \\ -E_{clu} \frac{xy}{r^3} & E_{clu} \frac{r^2 - y^2}{r^3} & -E_{clu} \frac{yz}{r^3} & 0 \\ -E_{clu} \frac{xz}{r^3} & -E_{clu} \frac{yz}{r^3} & E_{clu} \frac{r^2 - z^2}{r^3} & 0 \\ \frac{E}{E_{clu}} \cdot \frac{x}{r} & \frac{E}{E_{clu}} \cdot \frac{y}{r} & \frac{E}{E_{clu}} \cdot \frac{z}{r} & 1 \end{pmatrix}_{right}$$

- ErrorFlow:

$$\text{CovMat}(\vec{p}_{jet}, E_{jet}) = \sum_{PFO} \text{CovMat}(\vec{p}, E) \quad : \quad \sigma_{E_{jet}}^2 = \sigma_{conf}^2 + \sum_{PFO} \sigma_{E_{PFO}}^2$$

- MarlinKinfitProcessors:



$$\text{CovMat}(\vec{p}_{jet}, E_{jet}) \rightarrow (\sigma_{\theta_{jet}}, \sigma_{\phi_{jet}}, \sigma_{E_{jet}})$$



ErrorFlow: Jet Error Parametrisation from Particle Flow Objects (PFO)

Angles


The angular uncertainties obtained directly from track parameters / cluster position errors




⇒ Scale σ_θ and σ_ϕ by factor ~ 1.3 (for photons) and ~ 1.8 (for neutral hadrons)



Uncertainties in jet-level: θ & ϕ



Jet angular uncertainties need scaling factor ~ 1.6



Neutrino correction hypothesis

- ▶ Assign semi-leptonic decays to jets
- ▶ Add neutrino momentum to 4-momentum of assigned jet:

Test three hypothesis for neutrino energy in each semi-leptonic decay: E_ν^+ , E_ν^- , 0
 3^{nSLD} combination of E_ν 's for adding to jet 4-momentum:

Number of semileptonic decays in a jet: $nSLD = nSLDB + nSLDC$

Example:

If an event contains two jets: jet-1 contains 2 semi-leptonic decays and jet-2 contains 1 semi-leptonic decay,
27(= $3^2 \times 3^1$) combinations of E_ν 's are available for neutrino correction in the event:

▶ jet-1:

comb.	1	2	3	4	5	6	7	8	9
$\vec{p}_{\nu,1}$	-	+	0	-	+	0	-	+	0
$\vec{p}_{\nu,2}$	-	-	-	+	+	+	0	0	0

▶ jet-2:

comb.	1	2	3
$\vec{p}_{\nu,3}$	-	+	0

$\vec{p}_{\nu,1} + \vec{p}_{\nu,2}$ is added to 4-momentum of jet-1 and $\vec{p}_{\nu,3}$ is added to 4-momentum of jet-2.

$\vec{p}_{\nu,1} + \vec{p}_{\nu,2} + \vec{p}_{\nu,3} = 0$ allows fitter to neglect neutrino correction

Combination with highest fit probability is chosen as best neutrino correction.



Simple neutrino correction for Higgs mass reconstruction

► Neutrino energy correction:

Estimating neutrino energy as a fraction of corresponding lepton energy:

$$E_{jet}^{corr} = E_{jet} + E_\nu = E_{jet} + \left(\frac{1}{x} - 1\right) E_{lep}$$


► Uncertainty on jet energy parametrised as:

$$\sigma_{E_{jet}}^{corr} = \frac{100\%}{\sqrt{E_{jet}}} \oplus \sigma_\nu$$

$$\sigma_\nu^2 = \left(\frac{\sigma_{\langle x \rangle}}{\langle x \rangle^2}\right)^2 E_{lep}^2 + \left(\frac{1}{x} - 1\right) \Delta E_{lep}^2$$

► Fixed uncertainties on angles:

$$\Delta\theta_{jet} = \Delta\phi_{jet} = 100 \text{ mrad}$$




Blue: before neutrino energy correction

Orange: After neutrino energy correction

Simple correction to jet energy improves jet energy pull distribution as a measure of fit performance.



Simple neutrino correction for Higgs mass reconstruction



- Bias and assymetry in m_H is removed by correcting jet energy and adding ISR

Error flow and application in kinematic fit

Jet specific energy resolution for $e^+e^- \rightarrow ZH \rightarrow q\bar{q}b\bar{b}$ process at $\sqrt{s} = 350$ GeV

- ▶ Full 4×4 CovMatrix on 4-momentum of jets $\sigma(\vec{p}, E)$:
 - ▶ σ_{Det} : computed using subdetector momentum/energy resolution
 - ▶ σ_{Conf} : computed using jet energy and particle content (charged, neutral and photon)
 - ▶ $\sigma_\nu = 0.73 \cdot E_l$
 - ▶ $\sigma_{Had}, \sigma_{Clus}$ are not accounted for error flow procedure yet.
- ▶ Fixed (and wide) angular resolution: $\sigma_\theta = \sigma_\phi = 100$ mrad


Kinematic fit: vary jet quantities (E, θ, ϕ) within uncertainties $(\sigma_E, \sigma_\theta, \sigma_\phi)$

Improved fit probability by applying Error Flow on jet energy

⇒ Further improvements by error parametrization and handling sl-decays

DESY-THESIS-2017-045

$\sqrt{s} = 350$ GeV




fit constraints


momentum conservation: p_x

ISR is initialized to satisfy momentum conservation on x direction

- by error flow on jet energy



- by error flow on CovMatrix (new)



angular resolution for individual jets: improved constraint on momentum conservation



Reconstruction of b - and c - jets at e^+e^- Higgs Factories with ParticleFlow detectors | Yasser Radkhorrani | March 18, 2021 |




fit constraints


momentum conservation: p_y

ISR is initialized to satisfy momentum conservation on z direction

- by error flow on jet energy



- by error flow on CovMatrix (new)




Fit constraints


Momentum conservation: p_z

Adding 4-momentum of neutrino improves jet fit object initialization

- DBD 350/500 GeV samples



- MC-2020 250 GeV prod. samples



Proper neutrino correction for jets: improved constraint on momentum




fit constraints


energy conservation: E

Neutrino correction (best pre-fit \vec{p}_ν for successful fits) improves start values \Rightarrow better fit object initialization

- DBD 350/500 GeV samples



- MC-2020 250 GeV prod. samples



By neutrino correction, initial value of constraint function closer to target \Rightarrow fit should work better!

