

Contribution ID: 28 Type: not specified

Prospects of fast timing detectors for particle identificationat future Higgs factories

Tuesday 16 March 2021 11:00 (20 minutes)

Future e^+e^- colliders are excellent tools to probe fundamental physics beyond Standard Model via Higgs and electroweak precision measurements.

Modern silicon detectors are able to measure time-of-arrival with high precision of O(10 ps). This can be used to measure the time-of-flight (TOF) of the particles and improve their identification.

We develop reconstruction and calibration algorithms based on TOF information to separate π^{\pm} , K^{\pm} , p, \bar{p} particles at future Higgs factory detectors. Furthermore, we study how to implement fast timing silicon layers in the tracking and/or calorimeter systems, in order to derive requirements on the time resolution. As an example case, the ILD detector concept is studied.

The K^{\pm} mass measurement is a simple benchmark to test the performance of TOF algorithm. A precision at the level of 10 keV can be expected, which would significantly improve the knowledge of the K^{\pm} mass.

Time Zone

Europe/Africa/Middle East

Primary authors: Mr DUDAR, Bohdan (DESY); LIST, Jenny (Deutsches Elektronen-Synchrotron (DE)); ETE,

Remi (DESY)

Presenter: Mr DUDAR, Bohdan (DESY)

Session Classification: PD4/PD6: Software & Detector Performance / Calorimeters

Track Classification: Physics and Detectors Tracks: PD4: Software & Detector Performance