



Contribution ID: 101

Type: not specified

Di-Higgs production ($\gamma\gamma \rightarrow hh$) in Composite Models

Tuesday, 16 March 2021 09:00 (20 minutes)

In Standard Model (SM) Higgs Boson pair production initiated by photons ($\gamma\gamma \rightarrow hh$) is loop-generated process and thereby is very sensitive to any new couplings and particles that may come in loops. Composite Higgs Models (CHMs) provide an alternate mechanism to address the hierarchy problem of SM where Higgs could be a bound state of a strongly interacting sector instead of being an elementary field. These set of models apart from modifying the SM Higgs couplings could also introduce new effective couplings that can have substantial impact on the loop processes. In this work [1] we have studied the impact of Composite Higgs models in $\gamma\gamma \rightarrow hh$ (Di-Higgs) production process.

References

[1] A. Bharucha, G. Cacciapaglia, A. Deandrea, N. Gaur, D. Harada, F. Mahmoudi and K. Sridhar, [arXiv:2012.09470 [hep-ph]].

Time Zone

Europe/Africa/Middle East

Primary authors: DEANDREA, Aldo (Universit'e de Lyon, Universit'e Claude Bernard Lyon 1); BHARUCHA, Aoife (Aix Marseille Univ); Dr HARADA, Daisuke (Rudjer Boskovic Institute); MAHMOUDI, Farvah (Universit'e de Lyon, Universit'e Claude Bernard Lyon 1); CACCIAPAGLIA, Giacomo (Universit'e de Lyon, Universit'e Claude Bernard Lyon 1); SRIDHAR, K (Department of Theoretical Physics, Tata Institute of Fundamental Research); GAUR, Naveen (Department of Physics, Dyal Singh College (University of Delhi))

Presenter: Dr HARADA, Daisuke (Rudjer Boskovic Institute)

Session Classification: PD1/PD2: Theoretical Developments / Global Interpretations

Track Classification: Physics and Detectors Tracks: PD2: Global Interpretations