


# Exotica and Dark Matter searches

Michele Gallinaro  
LIP Lisbon  
April 28, 2021


- ✓ Introduction
- ✓ Dark matter
- ✓ Exotica searches

# 2012: A new boson discovery







# Standard Model theory of everything?

- Discovery of the Higgs boson marks the triumph of the SM
- However, even with the inclusion of the Higgs boson, SM is an incomplete theory



# Tests of the SM




# Beyond the Standard Model

The SM answers many of the questions about the structure of matter. But SM is not complete; still many unanswered questions:

- a) Why do we observe matter and almost no antimatter if we believe there is a symmetry between the two in the universe?
- b) What is this "dark matter" that we can't see that has visible gravitational effects in the cosmos?
- c) Are quarks and leptons actually fundamental, or made up of even more fundamental particles?
- d) Why are there three generations of quarks and leptons? What is the explanation for the observed pattern for particle masses?
- e) How does gravity fit into all of this?


# Dark matter and energy

- What is that accounts for 96% of the Universe?  
Nobody knows.
- It is one of the greatest mysteries of Science



# What can we look for?

A crowded field. At the LHC we can search for some of these



# How?

- Search for new phenomena
- Look for New Physics
- Indirect searches
  - precision measurements, event properties, etc.
- Direct searches
  - resonances, specific final states, model-(in)dependent searches, etc.
- Production and decay rates, event characteristics, advanced tools



# Dark Matter

## What is it?

- DM does not interact electromagnetically
- DM interacts gravitationally



Visual map

# Dark Matter (cont.)

Why is it interesting?


- We do not see it...but we feel it



# Dark Matter (cont.)


How do we find DM?

- Need to understand how it interacts with Universe
- Traditionally through a mediator
- Yields at least two new particles




# Searching for DM

## Particle Colliders



## Direct Detection




## Indirect Detection





# How do we find it: @underground

- Through a nuclear recoil




# How do we find it: @Space

- Through annihilation
  - Cosmic rays from DM




# How do we find it: @LHC

- Produced it through a mediator



# DM at the LHC


- CMS/ATLAS experiments **not** designed for DM searches



# DM searches at LHC

How do we find DM at the LHC?


- DM production gives MET signature



# DM searches: backgrounds

What are the backgrounds?


- $Z \rightarrow \nu\nu$ 
  - very similar to signal



# DM searches: backgrounds (cont.)

How to discriminate signal against the background?

- Look for high MET:



Study hadronic recoil

$$MET = -\sum_{\text{All particles}} p_T$$


$$MET(Z \rightarrow vv) = - Z \text{ recoil} + p_T(vv)$$

$$MET(Z \rightarrow vv) = - Z p_T$$


# DM searches: backgrounds (cont.)

How to discriminate signal against the background?


- Can fit the shape and look for signal



# DM+Z




- Main background is from ZZ di-bosons
- Understanding ZZ di-boson pT is critical



# Build a V-tagger


- Two jets are more collimated at high pT




- At low pT jets are “resolved”
  - Focus on reconstructing di-jets with mass near  $W$  mass
- At high pT get one “fat” jet
  - Focus on identifying one jet with mass near  $W$  mass
- Use additional variables to improve discrimination

# DM+jet/V


arXiv:1712.02345



## DM search in mono-jet/V



Need good control of systematics






# VBF: H(invisible)

arXiv:1809.06682, arXiv:1809.05937

- In the SM,  $B(H \rightarrow \text{invisible})$  only 0.1%
- Any significant deviation would indicate BSM
- Signature: Large MET,  $\Delta\phi(jj)$ , veto  $\ell/b$ -jets
  - C&C and shape fit of  $m(jj)$
- Main bkg: V+jets (95%)
- Tag with forward jets+MET
- Cross section  $\sim 4\text{pb}$
- Small background


Set limits:  $B(H \rightarrow \text{inv.}) < 0.37$  (0.28) @95%CL




# DM+Higgs

arXiv:1703.05236

- DM search with  $H \rightarrow b\bar{b}, \gamma\gamma$
- Model dependent search
- $Z'$  2HDM Model







- No significant excess
- Set limits for coupling  $g=0.8$



# Dark Matter



arXiv:1903.01400, arXiv:1901.0155, CMS-EXO-18-009

- Complementarity to direct/indirect searches
- **DM particles:**
  - interact via spin-0 & spin-1 mediators
  - are undetected (MET) recoiling against SM particle(s)
- Extensive program of mono-X searches ( $X = \text{jet}, \gamma, \text{lepton}, W, Z, t, tt, bb, H$ )
- No excess observed
- Interpretation through simplified models (DM and mediator masses, couplings to DM and SM)



# Experimental results


- Limits for given couplings between SM and DM interaction
- Competitive limits at low masses wrt other experiments



⇒ Collider results complement direct searches for low masses (<5-10 GeV)

# Resonant searches

## Resonant Searches




Among the highest dijet mass event recorded:  $m_{jj} = 8.12 \text{ TeV}$

$p_T = 3.8 \text{ TeV}$

$p_T = 3.8 \text{ TeV}$

# BSM models predict new resonances


- BSMs predict resonances with spin 0,1,2




- Are quarks fundamental particles?
  - Excited quarks in models of compositeness



- Randall-Sundrum (RS) models
  - Spin-2 graviton (KK-particle)



- Heavy-Vector Triplets
  - Spin-1 resonance
  - Models based on strength of vector boson interactions





- Sequential SM
  - $Z'$  and  $Z$  with same couplings to fermions
  - Width proportional to the mass


# New phenomena in di-jet events

CMS-EXO-17-026, arXiv:1910.08447

- Searches up to high masses
- QCD predicts a smooth, monotonic decrease in dijet invariant mass
- Search for a localized excess
- No significant excess observed




# Searching for dilepton resonances



CMS Experiment at the LHC, CERN

Data recorded: 2017-Jun-27 15:39:36.789504 GMT



Run / Event / LS: 297599 / 134277310 / 86



**Dimuon candidate event:  
Reconstructed mass of 2.4 TeV**

# Di-muon events



- Di-muon events: a re-discovery of the SM



# High-mass dilepton resonances


arXiv:1803.06292, arXiv:1903.06248, CMS-EXO-19-019

- Search for dilepton ( $ee, \mu\mu$ ) resonance




# Search for diboson resonances

- Heavy BSM resonances ( $>1\text{TeV}$ ) may decay into SM bosons (W,Z, H)
- Several final states
- Experimental challenges
  - SM bosons decay mostly to quarks
  - Due to large Lorentz boost, decay products merge into single jet
  - Clustered within a large-cone jet ( $R=0.8$ )
- Look into jet substructure
  - Jet “grooming”: get rid of soft jet components from UE/pileup, keep constituents from hard scatter
  - Apply filters (mass drop, pruning, trimming)



# Diboson resonances

- Many potential final states are possible
  - WW/WZ, ZW/ZZ, VV
- Hadronic channels with high sensitivity in high mass region




# Jet grooming


arXiv:0802.2470

## Mass drop/filtering


- Identify approx. symmetric sub-jets (with smaller mass than sum)



Mass drop: define 2 sub-jets



Filtering: re-cluster j1, j2 constituents




# Jet grooming (cont.)

arXiv:0912.1342, arXiv:0912.0033


## “Trimming”

- Uses  $k_T$  algorithm to make subjets (subjets with  $p_T^i/p_T^{\text{jet}} < f_{\text{cut}}$  removed)



## “Pruning”

- Recombine jet constituents, while veto wide-angle/softer constituents





# W, Z, H reconstruction

arXiv:1707.01303

- Grooming and jet mass

- Pruning
- soft drop (stable w/pileup, and good jet mass resolution  $\sim 10\%$ )


- Vector boson tagging ( $V \rightarrow q\bar{q}$ )
- $n$ -subjettiness  $\tau_{21}$ : how consistent with 2 sub-jets
- Categorization according to purity: high ( $< 0.35$ ) and high ( $> 0.35$ )



# $W, Z, H$ reconstruction (cont.)

arXiv:1707.01303



- Higgs boson tagging ( $H \rightarrow b\bar{b}$ )
  - Double b-tagging
  - Exploit b-tagging to identify two b-quarks in same jet
  - Soft-lepton information
  - Combines tracking and vertexing in MVA



# Searching for diboson resonances

arXiv:1906.08589




- No significant excess in any of the observed final states
- Exclusion limits: HVT models excluded up to 4.1 TeV, Spin-2 RS models up to 2.8 TeV
- Large improvements due to new methods for jet reconstructions and boson tagging



# Search for multi-lepton final states

JHEP 03(2020)051


- Search for new heavy particles
  - Heavy fermions/scalars produced in association with ttbar
- Search for 3 or more lepton final states
- Pair production of W/Z/H $\rightarrow\Sigma\Sigma$ 
  - Scalar sum of lepton  $p_T$  ( $L_T$ )
  - Bin and count ( $L_T$ +MET)





# X $\rightarrow$ VV $\rightarrow$ qqqq

arXiv:1708:05379

- All hadronic resonance search with single (qV) or double (VV) V-tag
  - At least 2 back-to-back jets  $p_T > 200\text{GeV}$
  - Categorization (jet mass,  $\tau_{21}$ )
- Background estimation: “bump hunt” fit data with power law





CMS Experiment at LHC, CERN  
Data recorded: Mon Jul 18 19:59:10 2016 CEST  
Run/Event: 276950 / 1080730125  
Lumi section: 573



# $X \rightarrow VH \rightarrow q\bar{q}bb$

arXiv:1707.01303

- All-hadronic search for  $V \rightarrow qq$  and  $H \rightarrow bb$  resonances
  - dedicated identification for  $H \rightarrow bb$  (b-tagging)
- Use categories
  - $V$ -jet mass ( $W$  or  $Z$ ),  $V$ -jet  $\tau_{21}$  (high-purity, low-purity),  $H$ -jet (tight and loose b-tag)






- Similar topology and background estimate to  $VV$  resonance search
- No significant excess found in data

# $X \rightarrow ZV \rightarrow llqq$

arXiv:1803.10093

- Search for resonances in  $Z \rightarrow ee/\mu\mu$ ,  $V \rightarrow qq$
- Clean final state (leptons)
  - Good mass resolution, good efficiency
- $\tau_{21}$  categorization (HP, LP)
- Parametrize main bkg ( $Z+jets$ ), fit to data in sidebands, take shape from MC





- Data compatible with SM-only hypothesis

# $X \rightarrow WV \rightarrow \ell\nu qq$

arXiv:1802.09407, B2G-19-002

- Search for a resonance decaying to WV in lepton+jet channel
- Categorization in  $\tau_{21}$  and W/Z mass
- Sideband+transfer function for bkg estimate



- Similar sensitivity to  $Z(\ell)\bar{V}(qq)$  search
- Excluded up to 1.1-3.1 TeV

# $X \rightarrow VH \rightarrow \ell\nu qq$


PLB 768(2017)137

- Search for a resonance decaying to VH in leptonic channels


- $Z \rightarrow \nu\nu$ : transverse mass  $m_T(VH)$
- $W \rightarrow \ell\nu$ : top control region
- $Z \rightarrow \ell\ell$ : high-efficiency dilepton ID
- $H(bb)$  b-tagging

- Sideband bkg prediction

- Heavy vector triplet ( $Z'$ ,  $W'$ )
- $g_V$ ,  $g_H$  ( $c_V$ ,  $c_F$ ): couplings




# Combination of diboson searches




# Combination of resonance searches

- Combination of searches for heavy resonances decaying to boson and lepton final states
- Large gain in statistical combination






# Resonance searches: Summary



# Leptoquarks

ATLAS-CONF-2021-008



- Pair production of leptoquarks
  - Study  $LQ \rightarrow b\tau$  final state
  - Can be interpreted in SUSY
- Main background from top quark pair events in different final states, determined from CRs
- Signal: use  $M_{T2}$  (stransverse mass) and  $s_T$  (scalar sum of  $\tau$  and jet pT) variables



# Heavy lepton resonances

EPJC 81(2021)218


- Two resonances in one process:  $W \rightarrow NL$
- Off-shell  $W$  decays to two new particles
- Signal selection:
  - OS and SS lepton pairs possible
  - two jets and MET
- Backgrounds from simulations and data CRs



# Vector-like quarks


## Motivation

- Simplest extension allowed in the quark sector
- Spin  $\frac{1}{2}$  fermions with vector coupling
- Can mix with SM quarks and modify their couplings to the W/Z/H bosons
- Sizeable mixing with 3<sup>rd</sup> family, b and t



## Properties



- Produced via strong and EWK interactions
- Mainly pair-produced
- Both CC and NC decay modes



# VLQ searches

PLB 779(2018)82



- Search for VLQ pair production decaying to WbWb
- Search in the **boosted regime**
- Can reconstruct the VLQ system



# Looking forward: PPS


CERN-LHC-2014-021

- The Precision Proton Spectrometer is a joint CMS and TOTEM project that aims at measuring the surviving scattered protons on both sides of CMS in standard running conditions
- Tracking and timing detectors inside the beam pipe at ~210m from IP5
- Project approved in Dec. 2014 by LHCC
- Data taking started in 2016 (full scope from 2017)



# PPS physics motivations


- Central Exclusive Production
  - photon-photon collisions
  - gluon-gluon fusion in color singlet,  $J^{PC}=0^+$
- High- $p_T$  system in central detector, together with very forward protons in PPS
  - momentum balance between central system and forward protons, provides strong kinematical constraints
  - Mass of central system measured by momentum loss of the two leading protons
- Gauge boson production by photon-photon fusion and anomalous couplings ( $\gamma\gamma WW$ ,  $\gamma\gamma ZZ$ , and  $\gamma\gamma\gamma$ )
- Search for new BSM resonances
- Study of QCD in a new domain



# Detectors



- Tracking detectors

- Goal: measure proton momentum
- Technology: silicon 3D pixels (6 planes per pot)




- Timing detectors

- Goal: identify primary vertex, reject “pileup”
- $\sigma_{\text{time}} \sim 10\text{ps} \Rightarrow \sigma_z \sim 2\text{mm}$
- Technology: silicon/diamond




# Roman Pot insertion

- Insertion procedure validated in 2016 by the LHC
  - Improvements carried out wrt earlier versions (RF shielding, cylindrical pots, ferrite, copper coating)
- Minimum distance of approach dramatically affects detector acceptance and physics reach
- A few mm ( $\sim 15\sigma$ ) from beam in nominal high-luminosity runs
  - Monitor beam losses, showers, interplay with collimators, beam impedance (heating, vacuum and beam orbit stability)





# LHC tunnel @ PPS location



# Timing detectors

Use timing to reject pileup background



- Two scenarios studied:
  - 10ps and 30ps time resolution
- Baseline: solid state detectors
- Detector options investigated:
  - Diamond sensors
  - Fast silicon sensors (UFSD, HFS)
- Status:
  - Diamond and LGAD detectors installed



# WW production

JHEP 08(2016)119


- Study of process:  $pp \rightarrow pWWp$ 
    - Clean process: W in central detector and “nothing” else, intact protons can be detected far away from IP
    - Exclusive production of W pairs via photon exchange: QED process, cross section well known
  - Backgrounds:
    - inclusive WW,  $\tau\tau$ , exclusive two-photon  $\gamma\gamma \rightarrow ll$ , etc.
  - Events:
    - WW pair in central detector, leading protons in PPS
  - SM observation of WW events
  - Anomalous coupling study
    - AQGCs predicted in BSM theories
    - parameters:  $a_0^W/\Lambda^2$ ,  $a_c^W/\Lambda^2$
  - Deviations from SM can be large




# Exclusive Dileptons

JHEP 07(2018)153


- Study exclusive processes at the EWK scale
- Search for two-photon production of opposite charge lepton pair with forward proton tagging



signal



Background: SD, DD, DY, dibosons, PU





- Signal selected with:
- at least one proton tagged, muons, kinematic selection

# Exclusive Dileptons (cont.)

JHEP 07(2018)153

- Correlation between the  $\xi$  values in central system vs RP
- 12 $\mu\mu$ , 8ee candidates observed ( $>5\sigma$  over expected bkg)
- First observation of two-photon production of a lepton pair at this mass range





# AQGC expected limits

arXiv:1607.03745

Expected limits @95%CL:

$$a_0^W/\Lambda^2 = 2 \times 10^{-6} (3 \times 10^{-6})$$




$$a_C^W/\Lambda^2 = 7 \times 10^{-6} (10 \times 10^{-6})$$



# $\gamma\gamma \rightarrow \gamma\gamma$ : Anomalous couplings, etc.


PRD 89(2014)114004, CMS-EXO-18-014

- Indirect search: neutral quartic gauge couplings (forbidden in SM) in  $\gamma\gamma \rightarrow \gamma\gamma$
- Expect to provide best sensitivity at LHC
- Sensitive to axion-like particles




# Prospects for Run3 and beyond

- More luminosity in a more challenging environment
- Will enhance the mass reach in the search for new particles
- Need to meet experimental challenges
  - Aging of detector, improve/adapt capability
  - Integrated luminosity: 300-3000/fb
  - peak luminosity of  $2 \times 10^{35} \text{ cm}^{-2} \text{s}^{-1}$
  - pileup will be ~150 or higher (Phase2)
  - large radiation doses



# BSM searches: resonances, etc.



# HL-LHC upgrades


Luminosity of  $\sim 3000 \text{ fb}^{-1}$  expected for HL-LHC

- Tracking information in “L1 track-trigger”
  - Tracker designed to enable finding all tracks  $w/p_T > 2\text{GeV}$  in  $< 4\mu\text{s}$
- Tracker is all silicon but with much higher granularity, up to  $|\eta|=4$ 
  - $> 2\text{billion pixels and strips}$
- High Granularity Endcap Calorimeters
  - Sampling of EM showers: every  $\sim 1\lambda$  (28 samples) w/pixels, and every  $\sim 0.35\lambda$  (24 samples) with pixels+scintillator to map 3D shower development
  - $\sim 6\text{M channels in all}$
- Precision timing to add a 4<sup>th</sup> dimension to object reconstruction

# Future: HL-LHC upgrades

## Trigger/HLT/DAQ

- Track information in hardware event selection
- 750 kHz hardware event selection
- 7.5 kHz events registered



## Barrel EM calorimeter

- New electronics
- Low operating temperature  $\approx -10^\circ$

## Muon systems

- New DT & CSC electronics
- New chambers  $1.6 < \eta < 2.4$
- Muon tagging  $2.4 < \eta < 3$

## New Endcap Calorimeters




- Rad. Tolerant
- 5D measurement

## New Tracker


- Rad. Tolerant - light
- High Definition measurement
- 40 MHz selective readout for hardware trigger
- Extended Pixel coverage to  $\eta \approx 3.8$

Beam radiation and luminosity  
Common systems and infrastructure

# Precision Timing Layer



- Time-of-flight precision  $\sim 30$ ps
  - $|\eta| < 3$ ,  $p_T > 0.7$ GeV
  - Crystal+SiPM: rad hard to  $2 \times 10^{14} n_{eq} \text{cm}^{-2}$
- Provide  $\sim \times 4\text{-}5$  effective PU reduction
  - 15% merged vertices reduce to 1.5%
  - Low PU track purity of vertices recovered
- Showers timed to 30ps in calorimeters



# Summary

- Excellent consistency of SM but **SM is incomplete**
- Direct and indirect searches for New Physics
  - Collected  $\sim 150/\text{fb}$  @13 TeV in 2015-2018
  - $\sim 300/\text{fb}$  to be collected in the next few years (up to LS3)
- Many studies performed with data collected so far
  - New dedicated algorithms being developed
  - Dark Matter, Exotica, signature-based searches
  - Other BSM searches
- Searches provide no hints for BSM yet

